





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/00		A2	(11) International Publication Number: WO 00/66095
			(43) International Publication Date: 9 November 2000 (09.11.00)
(21) International Application Number: PCT/GB00/01661		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 28 April 2000 (28.04.00)			
(30) Priority Data:			
9910166.9	30 April 1999 (30.04.99)	GB	
60/139,520	16 June 1999 (16.06.99)	US	
0002113.9	28 January 2000 (28.01.00)	GB	
(71) Applicant (for all designated States except US): STERIX LIMITED [GB/GB]; Magdalen Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): REED, Michael, John [GB/GB]; 42 Wimborne Gardens, London W13 8BZ (GB). POTTER, Barry, Victor, Lloyd [GB/GB]; 95 Dovers Park, Bathford, Bath BA1 7UE (GB).			
(74) Agents: ALCOCK, David et al.; D Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).			
		Published Without international search report and to be republished upon receipt of that report.	

(54) Title: **USE**

(57) Abstract

There is provided use of a cyclic compound or a pharmaceutically active salt thereof in the manufacture of a medicament to prevent and/or inhibit and/or arrest cell cycling, wherein the cyclic compound comprises at least one ring, wherein Group I and Group II, independently of each other, are attached to a ring of the cyclic compound; wherein Group I is a hydrocarbyl or an oxyhydrocarbyl group; and wherein Group II is a group of formula (I), X is P or S; when X is P, Y is =O or S, Z is -OH and R is hydrocarbyl or H; when X is S, Y is =O, Z is =O, and R is hydrocarbyl or N(R₁)(R₂), wherein each of R₁ and R₂ is independently selected from H or a hydrocarbyl group.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

USE

The present invention relates to a method and a use. In particular the present invention relates to a method for the manufacture of a medicament.

5

As discussed in "Molecular Cell Biology" 3rd Ed. Lodish *et al.* pages 177-181 different eukaryotic cells can grow and divide at quite different rates. Yeast cells, for example, can divide every 120 min., and the first divisions of fertilised eggs in the embryonic cells of sea urchins and insects take only 1530 min. because one large pre-existing cell is subdivided.

10 However, most growing plant and animal cells take 10-20 hours to double in number, and some duplicate at a much slower rate. Many cells in adults, such as nerve cells and striated muscle cells, do not divide at all; others, like the fibroblasts that assist in healing wounds, grow on demand but are otherwise quiescent.

15 Still, every eukaryotic cell that divides must be ready to donate equal genetic material to two daughter cells. DNA synthesis in eukaryotes does not occur throughout the cell division cycle but is restricted to a part of it before cell division.

The relationship between eukaryotic DNA synthesis and cell division has been thoroughly
20 analysed in cultures of mammalian cells that were all capable of growth and division. In contrast to bacteria, it was found, eukaryotic cells spend only a part of their time in DNA synthesis, and it is completed hours before cell division (mitosis). Thus a gap of time occurs after DNA synthesis and before cell division; another gap was found to occur after division and before the next round of DNA synthesis. This analysis led to the conclusion
25 that the eukaryotic cell cycle consists of an M (mitotic) phase, a G₁ phase (the first gap), the S (DNA synthesis) phase, a G₂ phase (the second gap), and back to M. The phases between mitoses (G₁, S, and G₂) are known collectively as the interphase.

Many nondividing cells in tissues (for example, all quiescent fibroblasts) suspend the cycle
30 after mitosis and just prior to DNA synthesis; such "resting" cells are said to have exited from the cell cycle and to be in the G₀ state.

It is possible to identify cells when they are in one of the three interphase stages of the cell

cycle, by using a fluorescence-activated cell sorter (FACS) to measure their relative DNA content: a cell that is in G_1 (before DNA synthesis) has a defined amount x of DNA; during S (DNA replication), it has between x and $2x$; and when in G_2 (or M), it has $2x$ of DNA.

5 The stages of mitosis and cytokinesis in an animal cell are as follows

(a) Interphase. The G_2 stage of interphase immediately precedes the beginning of mitosis. Chromosomal DNA has been replicated and bound to protein during the S phase, but chromosomes are not yet seen as distinct structures. The nucleolus is the only nuclear
10 substructure that is visible under light microscope. In a diploid cell before DNA replication there are two morphologic chromosomes of each type, and the cell is said to be $2n$. In G_2 , after DNA replication, the cell is $4n$. There are four copies of each chromosomal DNA. Since the sister chromosomes have not yet separated from each other, they are called sister chromatids.

15

b) Early prophase. Centrioles, each with a newly formed daughter centriole, begin moving toward opposite poles of the cell; the chromosomes can be seen as long threads. The nuclear membrane begins to disaggregate into small vesicles.

20 (c) Middle and late prophase. Chromosome condensation is completed; each visible chromosome structure is composed of two chromatids held together at their centromeres. Each chromatid contains one of the two newly replicated daughter DNA molecules. The microtubular spindle begins to radiate from the regions just adjacent to the centrioles, which are moving closer to their poles. Some spindle fibres reach from pole to pole; most
25 go to chromatids and attach at kinetochores.

(d) Metaphase. The chromosomes move toward the equator of the cell, where they become aligned in the equatorial plane. The sister chromatids have not yet separated.

30 (e) Anaphase. The two sister chromatids separate into independent chromosomes. Each contains a centromere that is linked by a spindle fibre to one pole, to which it moves. Thus one copy of each chromosome is donated to each daughter cell. Simultaneously, the cell elongates, as do the pole-to-pole spindles. Cytokinesis begins as the cleavage furrow

starts to form.

(f) Telophase. New membranes form around the daughter nuclei; the chromosomes uncoil and become less distinct, the nucleolus becomes visible again, and the nuclear membrane forms around each daughter nucleus. Cytokinesis is nearly complete, and the spindle disappears as the microtubules and other fibres depolymerise. Throughout mitosis the "daughter" centriole at each pole grows until it is full-length. At telophase the duplication of each of the original centrioles is completed, and new daughter centrioles will be generated during the next interphase.

10

(g) Interphase. Upon the completion of cytokinesis, the cell enters the G_1 phase of the cell cycle and proceeds again around the cycle.

It will be appreciated that cell cycling is an extremely important cell process. Deviations from normal cell cycling can result in a number of medical disorders. Increased and/or unrestricted cell cycling may result in cancer. Reduced cell cycling may result in degenerative conditions.

Cancer remains a major cause of mortality in most Western countries. Cancer therapies developed so far have included blocking the action or synthesis of hormones to inhibit the growth of hormone-dependent tumours. However, more aggressive chemotherapy is currently employed for the treatment of hormone-independent tumours.

Hence, the development of a pharmaceutical for anti-cancer treatment of hormone dependent and/or hormone independent tumours, yet lacking some or all of the side-effects associated with chemotherapy, would represent a major therapeutic advance.

The present invention seeks to provide a composition suitable for use in the treatment of cancers and, especially, breast cancer.

30

The following abbreviations used in the present specification have the given meanings:

E1S, oestrone sulphate;

2-MeOE2, 2-methoxyoestradiol;

2-OHE2, 2-hydroxyoestradiol;
EMATE, oestrone-3-*O*-sulphamate;
2-MeOEMATE, 2-methoxyoestrone-3-*O*-sulphamate;
2-MeOE1, 2-methoxyoestrone;
5 GenMATE, genistein-bis-sulphamate;
2-MeOE1S, 2-methoxyoestrone sulphate;
ER, estrogen receptor;
Tdt, terminal deoxynucleotidyl transferase;
TUNEL, Tdt-mediated dUTP-nick end labelling.

10

Oestrogens undergo a number of hydroxylation and conjugation reactions after their synthesis. Until recently it was thought that such reactions were part of a metabolic process that ultimately rendered oestrogens water soluble and enhanced their elimination from the body. It is now evident that some hydroxy metabolites (e.g. 2-hydroxy and 16 α -
15 hydroxy) and conjugates (e.g. oestrone sulphate, E1S) are important in determining some of the complex actions that oestrogens have in the body (1, 2).

20

Bradlow and his colleagues have investigated the formation of 2- and 16 α -hydroxylated oestrogens in relation to conditions that alter the risk of breast cancer. There is now evidence that factors which increase 2-hydroxylase activity are associated with a reduced cancer risk, while those increasing 16 α -hydroxylation may enhance the risk of breast cancer (3-6). Further interest in the biological role of estrogen metabolites has been stimulated by the growing body of evidence that 2-methoxyoestradiol (Fig 1, 1; 2-MeOE2) is an endogenous metabolite with anti-mitotic properties (7). 2-MeOE2 is formed from 2-
25 hydroxy estradiol (2-OHE2) by catechol estrogen methyl transferase, an enzyme that is widely distributed throughout the body.

30

Seegers and her colleagues originally reported that relatively high concentrations of 2-MeOE2 ($\geq 1\mu\text{M}$) were cytotoxic to MCF-7 breast cancer cells (8). They also observed that 2-MeOE2 caused uneven chromosome distribution in cells which also had a disorientated microtubule structure. A subsequent study revealed that 2-MeOE2 had no effect on the morphology of normal human skin fibroblasts, but a marked effect on transformed fibroblasts (9).

In vivo 2-MeOE2 inhibits the growth of tumours arising from the subcutaneous injection of Meth A sarcoma, B16 melanoma or MDA-MB-435 estrogen receptor negative (ER-) breast cancer cells (10, 11). It also inhibits endothelial cell proliferation and migration, and *in vitro* angiogenesis. It was suggested that the ability of 2-MeOE2 to inhibit tumour growth *in vivo* may be due to its ability to inhibit tumour-induced angiogenesis rather than direct inhibition of the proliferation of tumour cells (10).

The mechanism by which 2-MeOE2 exerts its potent anti-mitogenic and anti-angiogenic effects is still being elucidated. There is evidence that at high concentrations it can inhibit microtubule polymerisation and act as a weak inhibitor of colchicine binding to tubulin (12). Recently, however, at concentrations that block mitosis, tubulin filaments in cells were not found to be depolymerised but to have an identical morphology to that seen after taxol treatment (13). It is possible, therefore, that like taxol, a drug that is used for breast and ovarian breast cancer therapy, 2-MeOE2 acts by stabilising microtubule dynamics.

While the identification of 2-MeOE2 as a new therapy for cancer represents an important advance, the bioavailability of orally administered oestrogens is poor. Furthermore, they can undergo extensive metabolism during their first pass through the liver. As part of a research programme to develop a steroid sulphatase inhibitor for breast cancer therapy, oestrone-3-*O*-sulphamate (Fig 1, 2, EMATE) was identified as a potent active site-directed inhibitor (14, 15). Unexpectedly, EMATE proved to possess potent oestrogenic properties with its oral uterotrophic activity in rats being a 100-times higher than that of estradiol (16). Its enhanced oestrogenicity is thought to result from its absorption by red blood cells (rbcs) which protects it from inactivation during its passage through the liver and which act as a reservoir for its slow release for a prolonged period of time (17). A number of A-ring modified analogues were synthesised and tested, including 2-methoxyoestrone-3-*O*-sulphamate (3, 2-MeOEMATE) (18, 19). While this compound was equipotent with EMATE as a steroid sulphatase inhibitor, it was devoid of oestrogenicity.

The present invention seeks to provide a method of treatment of a cell cycling disorder. In particular, the present invention seeks to provide a method of treatment of cancer.

Aspects of the invention are defined in the appended claims.

The present invention is advantageous in that it provides a compound suitable for use in the treatment of cell cycling disorders such as cancers, including hormone dependent and
5 hormone independent cancers.

In addition, the present invention is advantageous in that it provides a compound that is suitable for use in the treatment of cancers such as breast cancer, ovarian cancer, endometrial cancer, sarcomas, melanomas, prostate cancer, pancreatic cancer etc. and
10 other solid tumours.

We have identified the effects that the compounds of the present invention such as 2-MeOEMATE, have on the proliferation of breast cancer cells and fibroblasts, and their effect on the cell cycle. In a preliminary *in vivo* study 2-MeOEMATE was found to cause
15 the rapid regression of nitrosomethylurea (NMU)-induced mammary tumours in intact rats.

In accordance with the present invention cell cycling is inhibited and/or prevented and/or arrested. Preferably cell cycling is prevented and/or arrested.


20 In a preferred aspect cell cycling is inhibited and/or prevented and/or arrested in the G₂/M phase.

In a preferred aspect cell cycling is irreversibly prevented and/or inhibited and/or arrested. Preferably cell cycling is irreversibly prevented and/or arrested.

25

By the term "irreversibly prevented and/or inhibited and/or arrested" it is meant after application of a compound of the present invention, on removal of the compound the effects of the compound, namely prevention and/or inhibition and/or arrest of cell cycling, are still observable. More particularly by the term "irreversibly prevented and/or inhibited and/or
30 arrested" it is meant that when assayed in accordance with Protocol I, cells treated with a compound of interest show less growth after Stage 2 of Protocol I than control cells. Protocol I is recited after the Examples.

The compound of the present invention is a cyclic compound comprising at least one ring, wherein Group I and Group II, independently of each other, are attached to a ring of the cyclic compound; wherein Group I is a hydrocarbyl or an oxyhydrocarbyl group; and wherein Group II is a group of the formula



X is P or S; when X is P, Y is =O or S, Z is -OH and R is hydrocarbyl or H; when X is S, Y is =O, Z is =O, and R is hydrocarbyl or N(R₁)(R₂), wherein each of R₁ and R₂ is independently selected from H or a hydrocarbyl group.

- 10 The term "hydrocarbyl group" as used herein means a group comprising at least C and H and may optionally comprise one or more other suitable substituents. Examples of such substituents may include halo-, alkoxy-, nitro-, a hydrocarbon group, an N-acyl group, a cyclic group etc. In addition to the possibility of the substituents being a cyclic group, a combination of substituents may form a cyclic group. If the hydrocarbyl group comprises
- 15 more than one C then those carbons need not necessarily be linked to each other. For example, at least two of the carbons may be linked *via* a suitable element or group. Thus, the hydrocarbyl group may contain hetero atoms. Suitable hetero atoms will be apparent to those skilled in the art and include, for instance, sulphur, nitrogen and oxygen.
- 20 In one preferred embodiment of the present invention, the hydrocarbyl group is a hydrocarbon group.

Here the term "hydrocarbon" means any one of an alkyl group, an alkenyl group, an alkynyl group, an acyl group, which groups may be linear, branched or cyclic, or an aryl

- 25 group. The term hydrocarbon also includes those groups but wherein they have been optionally substituted. If the hydrocarbon is a branched structure having substituent(s) thereon, then the substitution may be on either the hydrocarbon backbone or on the branch; alternatively the substitutions may be on the hydrocarbon backbone and on the branch.

- 30 The compound is a cyclic compound. In this regard, the compound can be a single ring

compound or a polycyclic compound. Here, the term "polycyclic" includes fused and non-fused ring structures including combinations thereof.

The cyclic group may be a single ring or it is a polycyclic ring structure.

5

In one aspect, the cyclic group may contain any one or more of C, H, O, N, P, halogen (including Cl, Br and I), S and P.

At least one of the cyclic groups may be a heterocyclic group (a heterocycle) or a non-heterocyclic group.

10

At least one of the cyclic groups may be a saturated ring structure or an unsaturated ring structure (such as an aryl group).

15 Preferably, at least one of the cyclic groups is an aryl ring.

Preferably, Group I and/or Group II is linked or attached to the aryl ring.

If the cyclic group is polycyclic some or all of the ring components of the compound may be fused together or joined *via* one or more suitable spacer groups.

20

The polycyclic compound may comprise a number of fused rings. In this aspect the fused rings may comprise any combination of different size rings, such as 3 six-membered rings (6,6,6), a six-membered ring, a seven-membered ring and a six-membered ring (6,7,6), a six-membered ring and two eight-membered rings (6,8,8) etc.

25

In one aspect the present invention relates to compounds wherein the polycyclic compounds are other than (6,6,7) rings. In a further aspect, the present invention relates to compounds wherein the polycyclic compounds only contain rings having other than 7 members.

30

In one aspect, if the cyclic group is polycyclic, Group I and Group II are each attached to the same ring of the polycyclic compound.

Thus, in accordance with one aspect of the present invention, preferably the compound is a polycyclic compound.


Preferably the polycyclic compound will contain, inclusive of all substituents, no more than
5 50 about carbon atoms, more usually no more than about 30 to 40 carbon atoms.

The polycyclic compound can comprise at least two ring components, or at least three ring components, or at least four ring components.

10 Preferably, the polycyclic compound comprises four ring components.

Preferred polycyclic compounds have a steroidal ring component - that is to say a cyclopentanophenanthrene skeleton, or bio-isosteres thereof.

15 As is well known in the art, a classical steroidal ring structure has the generic formula of:




In the above formula, the rings have been labelled in the conventional manner.

An example of a bio-isostere is when any one or more of rings A, B, C and D is a
20 heterocyclic ring and/or when any one or more of rings A, B, C and D has been substituted
and/or when any one or more of rings A, B, C and D has been modified; but wherein the
bio-isostere in the absence of the sulphamate group has steroidal properties.

In this regard, the structure of a preferred polycyclic compound can be presented as:

25



wherein each ring A', B', C' and D' independently represents a heterocyclic ring or a non-heterocyclic ring, which rings may be independently substituted or unsubstituted, saturated
5 or unsaturated.

By way of example, any one or more of rings A', B', C' and D' may be independently substituted with suitable groups - such as an alkyl group, an aryl group, a hydroxy group, a halo group, a hydrocarbyl group, an oxyhydrocarbyl group etc.

10

An example of D' is a five or six membered non-heterocyclic ring having at least one substituent.

In one preferred embodiment, the ring D' is substituted with a ethynyl group.

15

If any one of rings A', B', C' and D' is a heterocyclic ring, then preferably that heterocyclic ring comprises a combination of C atoms and at least one N atom and/or at least one O atom. Other heterocyclic atoms may be present in the ring.

20 Examples of suitable, preferred steroidal nuclei rings A'-D' of the compounds of the present invention include rings A-D of dehydroepiandrosterone and oestrogens including oestrone.

Preferred steroidal nuclei rings A'-D' of the compounds of the present invention include rings A-D of:

25

oestrones and substituted oestrones. viz:

oestrone

4-OH-oestrone

6 α -OH-oestrone

30 7 α -OH-oestrone

16 α -OH-oestrone

16 β -OH-oestrone

17-deoxyoestrone

oestrone

5

oestradiols and substituted oestradiols. viz:

4-OH-17 β -oestradiol

6 α -OH-17 β -oestradiol

7 α -OH-17 β -oestradiol

10 4-OH-17 α -oestradiol

6 α -OH-17 α -oestradiol

7 α -OH-17 α -oestradiol

16 α -OH-17 α -oestradiol

16 α -OH-17 β -oestradiol

15 16 β -OH-17 α -oestradiol

16 β -OH-17 β -oestradiol

17 α -oestradiol

17 β -oestradiol

17 α -ethinyl-17 β -oestradiol

20 17 β -ethinyl-17 α -oestradiol

17-deoxyoestradiol

oestriols and substituted oestriols. viz:

oestriol

25 4-OH-oestriol

6 α -OH-oestriol

7 α -OH-oestriol

17-deoxyoestriol

30 dehydroepiandrosterones and substituted dehydroepiandrosterones. viz:

dehydroepiandrosterones

6 α -OH-dehydroepiandrosterone

7 α -OH-dehydroepiandrosterone

16 α -OH-dehydroepiandrosterone

16 β -OH-dehydroepiandrosterone

- 5 In general terms the ring system A'B'C'D' may contain a variety of non-interfering substituents. In particular, the ring system A'B'C'D' may contain one or more hydroxy, alkyl especially lower (C₁-C₆) alkyl, e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, n-pentyl and other pentyl isomers, and n-hexyl and other hexyl isomers, alkoxy especially lower (C₁-C₆) alkoxy, e.g. methoxy, ethoxy, propoxy etc., alkynyl, e.g. ethynyl, or
10 halogen, e.g. fluoro substituents.

In an alternative embodiment, the polycyclic compound may not contain or be based on a steroid nucleus. In this regard, the polycyclic compound may contain or be based on a non-steroidal ring system - such as diethylstilboestrol, stilboestrol, coumarins, flavonoids,
15 combrestatin and other ring systems. Other suitable non-steroidal compounds for use in or as the composition of the present invention may be found in US-A-5567831.

Preferably, Group I and Group II are each attached to the same ring of the cyclic compound of the present invention at positions *ortho* with respect to each other.

20 Preferably, the polycyclic compound has a steroidal structure and Group I is attached to the A ring.

Preferably, the Group I is attached to the 2 position of the A ring of the steroidal structure.

25 Preferably, the polycyclic compound has a steroidal structure and Group II is attached to the A ring.

Preferably, the Group II is attached to the 3 position of the A ring of the steroidal structure.

30 Group I is a hydrocarbyl or an oxyhydrocarbyl group.

The term "hydrocarbyl group" as used herein is defined above.

In one preferred embodiment of the present invention, the hydrocarbyl group is a hydrocarbon group.

5 The term "hydrocarbon group" as used herein is defined above.

Preferably the hydrocarbyl group is of the formula C_{1-6} (such as a C_{1-3}).

10 If the compound comprises a steroidal nucleus, preferably the A ring has a hydrocarbyl group at the 2 position.

More preferably the group C_{1-6} is attached to the 2 position of the A ring of a steroidal nucleus.

15 Preferably, the hydrocarbyl group is an alkyl.

The alkyl is preferably a lower alkyl group containing from 1 to 5 carbon atoms, that is to say methyl, ethyl, propyl etc. Preferably, the alkyl group is methyl or ethyl.

20 Thus, in a preferred embodiment, if the compound comprises a steroidal nucleus the A ring has an methyl or ethyl substituent at the 2 position.

The term "oxyhydrocarbyl group" as used herein means a group comprising at least C, H and O and may optionally comprise one or more other suitable substituents. Examples of
25 such substituents may include halo-, alkoxy-, nitro-, an alkyl group, a cyclic group etc. In addition to the possibility of the substituents being a cyclic group, a combination of substituents may form a cyclic group. If the oxyhydrocarbyl group comprises more than one C then those carbons need not necessarily be linked to each other. For example, at least two of the carbons may be linked *via* a suitable element or group. Thus, the
30 oxyhydrocarbyl group may contain hetero atoms. Suitable hetero atoms will be apparent to those skilled in the art and include, for instance, sulphur and nitrogen.

In one preferred embodiment of the present invention, the oxyhydrocarbyl group is a

oxyhydrocarbon group.

Here the term "oxyhydrocarbon" means any one of an alkoxy group, an oxyalkenyl group, an oxyalkynyl group, which groups may be linear, branched or cyclic, or an oxyaryl group.

- 5 The term oxyhydrocarbon also includes those groups but wherein they have been optionally substituted. If the oxyhydrocarbon is a branched structure having substituent(s) thereon, then the substitution may be on either the hydrocarbon backbone or on the branch; alternatively the substitutions may be on the hydrocarbon backbone and on the branch.
- 10 Preferably the oxyhydrocarbyl group is of the formula $C_{1-6}O$ (such as a $C_{1-3}O$).

If the compound comprises a steroidal nucleus, preferably the A ring has an oxyhydrocarbyl group at the 2 position.

- 15 More preferably the group $C_{1-6}O$ is attached to the 2 position of the A ring of a steroidal nucleus.

Preferably, the oxyhydrocarbyl group is an alkoxy.

- 20 The alkyl group of the alkoxy substituent is preferably a lower alkyl group containing from 1 to 5 carbon atoms, that is to say methyl, ethyl, propyl etc. Preferably, the alkyl group is methyl.

- Thus, in a preferred embodiment, if the compound comprises a steroidal nucleus the A ring
25 has an methoxy substituent at the 2 position.

In one preferred embodiment of the present invention, preferably the compound is non-oestrogenic. The term "non-oestrogenic" means exhibiting no or substantially no oestrogenic activity.

30


In one preferred embodiment of the present invention, preferably the compound is not capable of being metabolised to compounds which display or induce hormonal activity.

In one preferred embodiment of the present invention, preferably the compound of the present invention is orally active.

The present invention is based on the highly surprising finding the compounds of the present invention provides an effective means to prevent and/or inhibit and/or arrest cell cycling.

We have found that compounds having a hydrocarbyl or an oxyhydrocarbyl substituent on the A ring and/or having a group of the formula of Group II on the A ring are potent (and in some cases highly potent) in preventing and/or inhibiting and/or arresting cell cycling.

A preferred compound of the present invention has the formula:




wherein rings A, B, C and D are independently optionally substituted.

Preferably Group I is in the 2-position.


Preferably Group II is in the 3-position.

In one preferred aspect, X is S and R is $N(R_1)(R_2)$. In other words, preferably Group II is a group of the formula:



wherein each of R_1 and R_2 is independently selected from H or a hydrocarbyl group.

Thus in a preferred aspect, the present invention provides the use of a cyclic compound or a pharmaceutically active salt thereof in the manufacture of a medicament to prevent and/or inhibit and/or arrest cell cycling, wherein the cyclic compound comprises at least one ring, wherein Group I and Group II, independently of each other, are attached to a ring of the cyclic compound; wherein Group I is a hydrocarbyl or an oxyhydrocarbyl group; and wherein Group II is a group of the formula



wherein each of R_1 and R_2 is independently selected from H or a hydrocarbyl group.

Group II of these preferred compounds is referred to as a "sulphamate group". These preferred compounds are referred to as "sulphamate compounds".

Preferably, R_1 and R_2 are independently selected from H or alkyl, cycloalkyl, alkenyl and aryl, or together represent alkylene, wherein the or each alkyl or cycloalkyl or alkenyl or optionally contain one or more hetero atoms or groups.

When substituted, the N-substituted sulphamate compound may contain one or two N-alkyl, N-alkenyl, N-cycloalkyl, N-acyl, or N-aryl substituents, preferably containing or each containing a maximum of 10 carbon atoms. When R_1 and/or R_2 is alkyl, the preferred values are those where R_1 and R_2 are each independently selected from lower alkyl groups containing from 1 to 5 carbon atoms, that is to say methyl, ethyl, propyl etc. Preferably R_1 and R_2 are both methyl. When R_1 and/or R_2 is aryl, typical values are phenyl and tolyl (-PhCH₃; *o*-, *m*- or *p*-). Where R_1 and R_2 represent cycloalkyl, typical values are cyclopropyl, cyclopentyl, cyclohexyl etc. When joined together R_1 and R_2 typically represent an alkylene group providing a chain of 4 to 6 carbon atoms, optionally interrupted by one or more hetero atoms or groups, e.g. -O- or -NH- to provide a 5-, 6- or 7- membered heterocycle, e.g.

morpholino, pyrrolidino or piperidino.

Within the values alkyl, cycloalkyl, alkenyl, acyl and aryl we include substituted groups containing as substituents therein one or more groups which do not interfere with the cell
5 cycling arresting and/or inhibiting and/or prevention activity of the compound in question. Exemplary non-interfering substituents include hydroxy, amino, halo, alkoxy, alkyl and aryl. A non-limiting example of a hydrocarbonyl group is an acyl group.

In some preferred embodiments, at least one of R_1 and R_2 is H.


10

Examples of suitable sulphonamide compounds for use in the present invention, or examples of suitable compounds that can be converted to suitable sulphonamide compounds for use in the present invention, can be found in the art - such as PCT/GB92/01587, PCT/GB97/03352, PCT/GB97/00444, GB 9725749.7, GB 9725750.5, US-A-5567831, US-
15 A-5677292, US-A-5567831, WO-A-96/05216, and WO-A-96/05217.

By way of example, PCT/GB92/01587 teaches novel sulphonamide compounds and pharmaceutical compositions containing them for use in the treatment of oestrogen dependent tumours, especially breast cancer. These sulphonamide compounds are sulphonamide esters.
20 Examples of such inhibitors are sulphonamide ester derivatives of steroids.

Another compound suitable for use in the present invention has at least the following skeletal structure:


25



wherein Group II denotes a sulphonamide group as described above; and rings A, B, C and D are independently optionally substituted.

Preferably, Group II is the above-mentioned preferred formula for the sulphamate group. In this regard, it is preferred that at least one of R_1 and R_2 is H.

Another compound suitable for use in the present invention has at least the following skeletal structure:




wherein Group II denotes a sulphamate group as described above; and
rings A, B, C and D are independently optionally substituted.

Preferably, R is the above-mentioned preferred formula for the sulphamate group. In this regard, it is preferred that at least one of R_1 and R_2 is H.

We have found that compounds having an a sulphamate group on the A ring and a hydrocarbyl or an oxyhydrocarbyl substituent on the A ring are particularly potent (and in some cases highly potent) in preventing and/or inhibiting and/or arresting cell cycling.

A preferred compound of the present invention has the formula:



Group II is a sulphamate group; and
rings A, B, C and D are independently optionally substituted.


Preferably Group I is in the 2-position.

Preferably Group II is in the 3-position.

5

For the present invention, preferably the sulphamate compound is an oxyhydrocarbyl steroidal sulphamate compound, in particular 2-methoxyoestrone-3-*O*-sulphamate, or a pharmaceutically active salt thereof, including analogues thereof.


- 10 2-methoxyoestrone-3-*O*-sulphamate is an analogue of oestrone-3-*O*-sulphamate (otherwise known as "EMATE"), which has the following structure:



- and can be called 2-methoxy EMATE.

- 15 2-methoxy EMATE is the sulphamoylated derivative of a naturally occurring oestrogen metabolite, 2-methoxyoestrone. This compound is formed in the liver by the hydroxylation of oestrone by a 2-hydroxylase, with subsequent metabolism to the methoxy derivative by catechol oestrogen methyl transferase.

- 20 2-methoxy EMATE has the formula presented as formula below:



2-methoxy EMATE is believed to act *in vivo*, at least in part, by preventing and/or

inhibiting and/or arresting cell cycling.

Thus, in a highly preferred embodiment the sulphamate compound is an oxyhydrocarbyl steroidal sulphamate compound, in particular 2-methoxyoestrone-3-*O*-sulphamate (2-methoxy EMATE).

In this regard, we have found that a sulphamate compound having a C₁₋₆ (such as a C₁₋₃) alkoxy substituent at the 2 position of the A ring, in particular 2-methoxy EMATE, is highly potent in preventing and/or inhibiting growth of tumours.

In one embodiment, preferably, the sulphamate compound is an oxyhydrocarbyl steroidal sulphamate compound wherein the sulphamate group is in the 3 position on the steroidal component and/or the oxyhydrocarbyl group is in the 2-position position on the steroidal component.

In one embodiment, preferably, the sulphamate compound is an oxyhydrocarbyl derivative of oestrone sulphamate.

In one embodiment, preferably, the sulphamate compound is an oxyhydrocarbyl derivative of oestrone-3-*O*-sulphamate.

In one embodiment, preferably, the sulphamate compound is a C₁₋₆ (such as a C₁₋₃) alkoxy derivative of oestrone-3-*O*-sulphamate.


In one embodiment, preferably, the sulphamate compound is a 2-C₁₋₆ (such as a C₁₋₃) alkoxy derivative of oestrone-3-*O*-sulphamate.

In one embodiment, preferably, the sulphamate compound is 2-methoxyoestrone-3-*O*-sulphamate.

For the present invention, the sulphamate compound may preferably be a hydrocarbyl steroidal sulphamate compound, in particular 2-ethyloestrone-3-*O*-sulphamate, or a pharmaceutically active salt thereof, including analogues thereof.

2-ethyloestrone-3-*O*-sulphamate is an analogue of oestrone-3-*O*-sulphamate and can be called 2-ethyl EMATE.

- 5 2-ethyl EMATE has the formula presented as formula below:



- 10 2-ethyl EMATE is believed to act *in vivo*, at least in part, by preventing and/or inhibiting and/or arresting cell cycling.

Thus, in a highly preferred embodiment the sulphamate compound is a hydrocarbyl steroidal sulphamate compound, in particular 2-ethyloestrone-3-*O*-sulphamate (2-ethyl EMATE).

- 15 In this regard, we have found that a sulphamate compound having a C₁₋₆ (such as a C₁₋₃) alkyl substituent at the 2 position of the A ring, in particular 2-ethyl EMATE, is highly potent in preventing and/or inhibiting growth of tumours.

- 20 In one embodiment, preferably, the sulphamate compound is a hydrocarbyl steroidal sulphamate compound wherein the sulphamate group is in the 3 position on the steroidal component and/or the hydrocarbyl group is in the 2-position position on the steroidal component.

- 25 In one embodiment, preferably, the sulphamate compound is a hydrocarbyl derivative of oestrone sulphamate.

In one embodiment, preferably, the sulphamate compound is a hydrocarbyl derivative of oestrone-3-*O*-sulphamate.

In one embodiment, preferably, the sulphamate compound is a C₁₋₆ (such as a C₁₋₃) alkyl derivative of oestrone-3-*O*-sulphamate.


- 5 In one embodiment, preferably, the sulphamate compound is a 2-C₁₋₆ (such as a C₁₋₃) alkyl derivative of oestrone-3-*O*-sulphamate.

In one embodiment, preferably, the sulphamate compound is 2-ethyloestrone-3-*O*-sulphamate.

10

The sulphamate compounds of the present invention may be prepared by reacting an appropriate alcohol with the appropriate sulphonyl chloride, R₁R₂NSO₂Cl. Preferred conditions for carrying out the reaction are as follows. Sodium hydride and a sulphonyl chloride are added to a stirred solution of the alcohol in anhydrous dimethyl formamide at 15 0°C. Subsequently, the reaction is allowed to warm to room temperature whereupon stirring is continued for a further 24 hours. The reaction mixture is poured onto a cold saturated solution of sodium bicarbonate and the resulting aqueous phase is extracted with dichloromethane. The combined organic extracts are dried over anhydrous MgSO₄. Filtration followed by solvent evaporation *in vacuo* and co-evaporated with toluene affords a 20 crude residue which is further purified by flash chromatography. Preferably, the alcohol is derivatised, as appropriate, prior to reaction with the sulphonyl chloride. Where necessary, functional groups in the alcohol may be protected in known manner and the protecting group or groups removed at the end of the reaction.

- 25 In one aspect of the present invention X is P, Y is =O and Z is -OH; or X is S; Y is =O, Z is =O, and R is hydrocarbyl. Alternatively stated, the present compound is a sulphonate or a phosphonate compound in which Group I is a group of the formula:



wherein X is P or S; when X is P, Z is -OH; when X is S, Z is =O; and R is hydrocarbyl.


30

Preferably the sulphonate or phosphonate compound is a polycyclic compound. More preferably the sulphonate or phosphonate compound has a steroidal structure.

When the sulphonate or phosphonate compound has a steroidal structure the sulphonate or phosphonate compound may have at least one sulphonate or phosphonate group attached to the 3 position of the A ring of the steroidal nucleus.

In a further preferred aspect the sulphonate or phosphonate compound comprises at least one hydrocarbon group, preferably C₁₋₆ alkyl, attached to the 2 position of the A ring of a steroidal nucleus.

Thus in a further aspect the present invention provides a sulphonate or a phosphonate compound comprising a steroidal ring and a sulphonate or a phosphonate group of the formula:



wherein

X is P or S;

when X is P, Z is -OH, and R is hydrocarbyl or H;

when X is S, Z is =O, and R is hydrocarbyl; and

wherein the sulphonate or phosphonate group is attached to the 3 position of the A ring of the steroidal nucleus; and at least one hydrocarbon group, preferably C₁₋₆ alkyl, is attached to the 2 position of the A ring of a steroidal nucleus.

These novel sulphonate or a phosphonate compounds may be used in medicine, preferably used in accordance with the present invention.

Examples of suitable sulphonate or a phosphonate compounds for use in the present invention, or examples of suitable compounds that can be converted to suitable sulphonate or a phosphonate compounds for use in or the present invention, can be found in the art - such as PCT/GB92/01586.

In a further aspect the present invention provides the use of a composition in the manufacture of a medicament to prevent and/or inhibit and/or arrest cell cycling, wherein the composition comprises

- 5 (i) a compound as defined herein; and
- (ii) a further compound as defined herein; and/or
- (iii) a pharmaceutically acceptable carrier, diluent, or excipient; and/or
- (iv) a biological response modifier; and/or
- (v) pure antioestrogens; and/or
- 10 (vi) selective estrogen response modifiers (SERMs); and/or
- (vii) taxol.

The term biological response modifier ("BRM") includes cytokines, immune modulators, growth factors, haematopoiesis regulating factors, colony stimulating factors, chemotactic,
15 haemolytic and thrombolytic factors, cell surface receptors, ligands, leukocyte adhesion molecules, monoclonal antibodies, preventative and therapeutic vaccines, hormones, extracellular matrix components, fibronectin, etc.

Preferably, the biological response modifier is a cytokine. Examples of cytokines include:
20 interleukins (IL) - such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-19; Tumour Necrosis Factor (TNF) - such as TNF- α ; Interferon alpha, beta and gamma; TGF- β .

Preferably the cytokine is tumour necrosis factor (TNF).

25

For the preferred aspect of the present invention, the TNF may be any type of TNF - such as TNF- α , TNF- β , including derivatives or mixtures thereof. More preferably the cytokine is TNF- α .

30 Teachings on TNF may be found in the art - such as WO-A-98/08870 and WO-A-98/13348.

For pharmaceutical administration, the composition of the present invention can be

formulated in any suitable manner utilising conventional pharmaceutical formulating techniques and pharmaceutical carriers, adjuvants, excipients, diluents etc. - such as those for parenteral administration. Approximate effective dose rates are in the range 100 to 800 mg/day depending on the individual activities of the compounds in question and for a patient of average (70Kg) bodyweight. More usual dosage rates for the preferred and more active compositions will be in the range 200 to 800 mg/day, more preferably, 200 to 500 mg/day, most preferably from 200 to 250 mg/day. They may be given in single dose regimes, split dose regimes and/or in multiple dose regimes lasting over several days. For oral administration they may be formulated in tablets, capsules, solution or suspension containing from 100 to 500 mg of composition per unit dose. Alternatively and preferably the compositions will be formulated for parenteral administration in a suitable parenterally administrable carrier and providing single daily dosage rates in the range 200 to 800 mg, preferably 200 to 500, more preferably 200 to 250 mg. Such effective daily doses will, however, vary depending on inherent activity of the active ingredient and on the bodyweight of the patient, such variations being within the skill and judgement of the physician.

The compound or composition of the present invention may be administered in any suitable manner - such as any one or more of oral administration, topical administration (such as by means of a patch), parenteral administration, rectal administration or by inhalation spray.

In the method of treatment, the subject is preferably a mammal, more preferably a human. For some applications, preferably the human is a woman.

For particular applications, it is envisaged that the compounds or compositions of the present invention may be used in combination therapies, either with a sulphatase inhibitor, or, for example, in combination with an aromatase inhibitor, such as for example, 4-hydroxyandrostenedione (4-OHA).


In accordance with the present invention, the components of the composition can be added in admixture, simultaneously or sequentially. Furthermore, in accordance with the present invention it may be possible to form at least a part of the composition *in situ* (such as *in vivo*) by inducing the expression of - or increasing the expression of - one of the components. For example, it may be possible to induce the expression of - or increase the

expression of - the biological response modifier, such as TNF. By way of example, it may be possible to induce the expression of - or increase the expression of - TNF by adding bacterial lipopolysaccharide (LPS) and muramyl dipeptide (MDP). In this regard, bacterial LPS and MDP in combination can stimulate TNF production from murine spleen cells *in vitro* and tumour regression *in vivo* (Fuks *et al* Biull Eksp Biol Med 1987 104: 497-499).

The present invention also provides compositions/compounds which:

- cause inhibition of growth of oestrogen receptor positive (ER+) and ER negative (ER-) breast cancer cells *in vitro* by preventing and/or inhibiting and/or arresting cell cycling.
- cause regression of nitroso-methyl urea (NMU)-induced mammary tumours in intact animals (i.e. not ovariectomised).
- prevent and/or inhibit and/or arrest cell cycling in cancer cells.
- act *in vivo* by preventing and/or inhibiting and/or arresting cell cycling
- act as a cell cycling agonist.

In a further aspect the present invention provides use of a cyclic compound or a pharmaceutically active salt thereof in the manufacture of a cell cycling agonist, wherein the cyclic compound comprises at least one ring, wherein Group I and Group II, independently of each other, are attached to a ring of the cyclic compound; wherein Group I is a hydrocarbyl or an oxyhydrocarbyl group; and wherein Group II is a group of the formula



X is P or S; when X is P, Y is =O or S, Z is -OH and R is hydrocarbyl or H; when X is S, Y is =O, Z is =O, and R is hydrocarbyl or N(R₁)(R₂), wherein each of R₁ and R₂ is independently selected from H or a hydrocarbyl group.

The compounds of the present invention such as 2-methoxyoestrogens are emerging as a new class of drug that can inhibit tumour growth and inhibit angiogenesis. As sulphamoylation of oestrogens enhances their potency and bioavailability we describe

- below the synthesis of 2-methoxyoestrone-3-*O*-sulphamate (2-MeOEMATE) and comparison of its ability to inhibit the proliferation of breast cancer cells with that of 2-methoxyoestrone (2-MeOE1). 2-MeOEMATE (1 μ M) inhibited the growth of estrogen receptor positive MCF-7 breast cancer cells by 52% whereas 2-MeOE1 had little effect at this concentration. 2-MeOEMATE also inhibited the growth of estrogen receptor positive MDA-MB-231 breast cancer cells. Exposure of cells to 2-MeOEMATE caused them to round up and become detached suggesting that this compound may induce cells to undergo apoptosis. Cell cycle analysis revealed that 2-MeOEMATE caused cells to arrest in the G₂/M phase with the increase in G₂/M arrested cells being detectable by 12h. Exposure of MCF-7 cells to 2MeOEMATE for 24h followed by culture in drug-free medium for 24h did not reverse the arrest of cells in the G₂/M phase. TUNEL analysis confirmed that 2-MeOEMATE induced apoptosis in a significant proportion of treated MCF-7 cells. In an preliminary *in vivo* study, employing nitrosomethylurea-induced mammary tumours in intact rats, 2-MeOE1 (20mg/kg/d, p.o. for 11 days) had little effect on tumour growth. In contrast, the same dose of 2-MeOEMATE resulted in the almost complete regression of 2/3 tumours over an 11-day period. It is concluded that 2-MeOEMATE should have considerable therapeutic potential for the treatment of hormone-dependent and independent breast tumours.
- In summation, the present invention provides compositions for use in treatment of tumours and pharmaceutical compositions containing them.

The present invention will now be described only by way of example, in which reference, in which reference shall be made to the following Figures.

- Figure 1 which illustrates structures;
Figure 2 which is a graph;
Figure 3 which is a photographic plate;
Figure 4 which is a photographic plate
Figure 5 which is a photographic plate
Figure 6 which is a graph;
Figure 7 which is a graph;
Figure 8 which is a graph;

Figure 9 which is a graph;

Figure 10 which is a graph;

Figure 11 which is a graph;

Figure 12 which illustrates structures;

5 Figure 13 which are graphs;

Figure 14 which are graphs;

Figure 15 which is a photographic plate; and

Figure 16 which are graphs.

10 Figure 1 Structures: compound 1, 2-methoxyoestradiol (2-MeOE2); 2, oestrone-3-*O*-sulphamate (EMATE); 3, 2-methoxyoestrone-3-*O*-sulphamate (2-MeOEMATE); 4, 2-methoxyoestrone (2-MeOE1); 5, genistein bis-sulphamate (GenMATE); 6, genistein, 7, 2-methoxyoestrone-3-sulphate (2-MeOE1S).

15 Figure 2 Dose-response showing the effects of 2-methoxyoestrone (2-MeOE1) or 2-methoxyoestrone-3-*O*-sulphamate (2-MeOEMATE) on the proliferation of MCF-7 breast cancer cells. Cells (5000 per well) were exposed to drug for 4 days (means \pm s.d., $n=3$).

Figure 3 Effect of a) vehicle, b) 2-methoxyoestrone-3-*O*-sulphamate (2-MeOEMATE, 1 μ M) or 2-methoxyoestrone (2-MeOE1, 1 μ M) on MCF-7 breast cancer cells after 24h.

Figure 4 Effect of a) vehicle or b) 2-methoxyoestrone-3-*O*-sulphamate (2-MeOEMATE, 1 μ M) on MDA-MB-231 breast cancer cells after 24h.

25

Figure 5 Effect of a) vehicle, b) 2-methoxyoestrone-3-*O*-sulphamate (2-MeOEMATE, 1 μ M) or c) 2-MeOEMATE (5 μ M) on breast tumour-derived fibroblasts at 24h.

30 Figure 6 DNA histograms of vehicle (A) or 2-methoxyoestrone-3-*O*-sulphamate (10 μ M) treated MCF-7 breast cancer cells at 4h (B), 12h (C) or 24h (D). By 12 h after treatment there was evidence of an arrest of cells at the G₂/M phase.

Figure 7 DNA histogram of 2-methoxyoestrone-3-*O*-sulphamate (10 μ M) treated MCF-7 breast cancer cell s. Compared with controls at 24h (A) or 48h (C) cells treated with drug showed a significant increase in the proportion of cells in the G₂/M phase at 24h (B) and 48h (D), but also an increase in the sub-G₁ fraction. After exposure of cells for 24h, removal of the drug with a further 24h culture in drug-free medium did not reverse the arrest of cells in the G₂/M phase (E).

Figure 8 TUNEL analysis of control MCF-7 cells (A) or cells exposed to 2-methoxyoestrone-3-*O*-sulphamate (10 μ M) for 48h (B). Histograms are overlays for cells stained in the absence of TdT (bold) or cells stained with TUNEL reaction mixture (open). The bar in the lower figure represents TUNEL positive cells and represents approximately 10% of treated cells.

Figure 9 Effect of vehicle (propylene glycol, control animals 1-3), 2-methoxyoestrone (2-MeOE1, 20mg/kg/d, p.o. for 11 days in two animals) or 2-methoxyoestrone-3-*O*-sulphamate (20mg/kg/d, p.o. for 11 days in 3 animals) on the growth of nitrosomethylurea-induced mammary tumours in intact rats. Results are expressed as the tumour volumes (expressed as a percentage) on days 6 or 11 (Vdn) compared with their volumes at the start of drug administration.

Figure 10 Effect of 2-methoxy or 2-ethyl oestrone or their sulphamate derivatives on growth of MCF-7 breast cancer cells.

Figure 11 Irreversible effect of 2-methoxy or 2-ethyl oestrone sulphamate on growth of MCF-7 breast cancer cells after prior exposure.

Figure 12. Structures: compound 8, 2-Acetyloestrone 3-methyl ether; 9, 2-Ethylestrone 3-methyl ether; 10, 2-Ethylestrone; 11, 2-Ethylestrone 3-*O*-sulphamate.

Figure 13 Effects of 2-EtEMATE on the DNA content of MCF7, ZR-75-1, CAL51 and CAMA1 cells using flow cytometry of propidium iodide (PI) stained cells.

Figure 14 Effects of 2-MeOE2, 2-MeOE1, 2-EtE1, 2-MeOEMATE, and 2-EtEMATE,

on MCF7, ZR-75-1, CAL51 and CAMA1.

Figure 15 Photographic plate of stained nuclei after 2EtEMATE exposure.

5 Figure 16 Effects of 2-EtEMATE, on MCF7, ZR-75-1, CAL51 and CAMA1.

General Methods

10 All reagents and solvents employed were of general purpose or analytical grade unless otherwise stated, and purchased from either Aldrich or Sigma Chemicals or Lancaster Synthesis.

Silica gel refers to silica gel, Merck, grade 60. Product(s) and starting material were
15 detected either viewing under UV light or treating with a methanolic solution of phosphomolybic acid followed by heating. NMR spectra were determined using acetone-
 d_6 , $CDCl_3$ or $DMSO-d_6$ as solvent and TMS as internal standard, unless otherwise stated. The 1H NMR and ^{13}C NMR spectra were recorded on a Jeol GX 270 at 270 MHz and on a
Jeol EX 400 at 400 MHz NMR spectrometer. The following abbreviations are used to
20 describe resonances in 1H NMR and ^{13}C NMR spectra: s, singlet; d, doublet; br, broad; t, triplet; q, quartet; m, multiplet and combination such as dd, doublet of doublets. IR spectra
were determined as KBr discs, using a Perkin-Elmer 782 Infra-Red Spectrophotometer. Melting points were determined on a Reichert-Jung Kofler Block and are uncorrected.
Mass spectra were recorded on VG 7070 and VG Autospec instruments at the Mass
25 Spectrometry Service at the University of Bath. FAB-mass spectra were carried out using *m*-nitrobenzyl alcohol (*m*-NBA) as the matrix. HPLC stability studies were determined
using LDC Constametric 3000 HPLC Pump and Spectrometer 3000 variable wave length detector. CHN analysis was determined using gas chromatography at the Microanalysis
Service at the University of Bath.

30

All reagents and solvents used were stored away from moisture and light and dried before use. Low temperature experiments were conducted using a well insulated external bath containing either ice/water with NaCl for 0°C or carbon dioxide pellets with acetone or

using cold plate. Experiments requiring anhydrous conditions were guarded by mean of a drying tube containing self-indicating silica. Evaporation of solvents was carried out with a rotary evaporator at reduced pressure (water pump) and on stated occasions, followed by the use of a high vacuum pump. Samples were dried in drying tube under high vacuum and
5 low temperature.

All assays were performed at the Department of Endocrinology and Metabolic Medicine, Imperial College School of Medicine, St. Mary's Hospital, London.

10 Preparation of sulphamoyl chloride

Sulphamoyl chloride was prepared by the reaction of chlorosulphonyl isocyanate with formic acid according to the method of Appel and Berger.⁴⁴ To anhydrous sulphur-free toluene (150 ml) chlorosulphonyl isocyanate (25 g., 177 mmol) was added at 0°C under an
15 atmosphere of N₂. After stirring, formic acid (6.0 ml, 156 mmol) was added dropwise at 0°C under N₂. The resulting white light emulsion was kept stirring overnight and the toluene removed by using a water vacuum pump to give a light yellow crude of sulphamoyl chloride (16.24 g, 79%). A standard solution (ca 0.70 M) of sulphamoyl chloride was then prepared by dissolving the crude crystalline product in anhydrous sulphur-impurities-free
20 toluene and stored in the refrigerator under N₂. No titration was attempted on this sulphamoyl chloride solution whose molarity was estimated according to the weight of the original crude sulphamoyl chloride obtained after workup. Toluene used for preparing sulphamoyl chloride solution was purified according to the method described.⁴⁵ Cold toluene (1-3 litres) was placed in a separating funnel and washed with cold conc. H₂SO₄
25 (100 ml/litres, 3-4 times), once with water, once with aqueous 5% NaOH and again with water until neutral, dried with anhydrous MgSO₄ followed by sodium metal overnight and then fractionally distilled under N₂ from the sodium metal and stored in dark under N₂.

Anhydrous formic acid used for preparing sulphamoyl chloride was purified according the
30 method described.⁴⁵ Formic acid (98%) was stirred overnight with boric anhydride and then distilled under N₂, stored in dark under N₂.

General method for sulphamoylation

Starting with the parent compound, the sulphamate derivatives were prepared essentially as described by Howarth *et al.*¹⁴ unless stated otherwise. In this regard, a solution of the appropriate parent compound in anhydrous DMF was treated with sodium hydride [60% dispersion; 1.2 and 2.5 equiv. for monohydroxyl and dihydroxyl compounds respectively, unless stated otherwise] at 0°C under an atmosphere of N₂. After evolution of hydrogen had ceased, sulphamoyl chloride in toluene [excess, ca. 5-6 eq.] was added and the reaction mixture was poured into brine after warming to room temperature overnight and ethyl acetate was added. The organic fraction was washed exhaustively with brine, dried (MgSO₄), filtered and evaporated under reduced pressure. The crude product obtained was purified by flash chromatography or preparative TLC followed by recrystallisation to give the corresponding sulphamate. All the compounds were characterised by spectroscopic and combustion analysis.

Synthesis of 2-methoxyoestrone-3-*O*-sulphamate (2-methoxy EMATE)

2-methoxy EMATE (Fig 1, 3, 2-MeOEMATE) was synthesised by treating a solution of 2-methoxyoestrone (Fig 1, 4, 2-MeOE1) in anhydrous dimethylformamide with sodium hydride at 0°C. After evolution of hydrogen had ceased sulphamoyl chloride (2 equiv.) was added and the reaction mixture was allowed to warm to room temperature overnight. The compound was purified by silica gel flash chromatography, was a single pure spot by TLC and exhibited satisfactory spectroscopic and microanalytical data.

In this regard, 2-Methoxy oestrone (75 mg, 0.250 mmol) gave a crude product (103 mg) which was fractionated on silica (50 g) with chloroform/acetone (8:1) and upon evaporation the second fraction gave a pale white residue (83 mg, 81%) which was recrystallised in ethylacetate/hexane (1:2) to give 1 as white crystals (69 mg) .m.p = 177-180°C. R_fs= 0.29 and 0.54 for chloroform/ acetone 8:1 and 4:1 respectively and 0.46 and 0.31 for ethylacetate/hexane 2:1 and 1:1 respectively. ν_{max} (KBr) 3400, 3300 (-NH₂), 1610 (C=O), and 1380 (-SO₂N-) cm⁻¹. δ_{H} (CDCl₃) 0.922 (3H, s, C-18-CH₃), 1.24- 2.87 (15H, m), 3.88 (3H, s, C-2-OCH₃), 5.0 (2H, br s, exchanged with D₂O, - SO₂NH₂), 6.93 (1H, s, C-1- H) and 7.06 (1H, s, C-4-H). MS: m/z (+ve ion FAB in m- NBA, rel.

intensity) 379.1 [100, (M)⁺], 300.0 [25, (M-SO₂NH₂)⁺]. MS: *m/z* (-ve ion FAB in *m*-NBA, rel. intensity) 378.0 [100, (M-H)⁺]. Acc. MS: *m/z* (FAB⁺) = 380.1515. C₁₉H₂₆NO₅S requires 380.1532 Found C, 60.0; H, 6.7; N, 3.67; C₁₉H₂₅NO₅S requires C, 60.14; H, 6.64; N, 3.69%.

5

Genistein bis-sulphamate (Fig 1, 5, GenMATE) was prepared from genistein (Fig 1, 6) by the same procedure with the exception that 2.5 eq sodium hydride was used.

2-Methoxyestrone and other steroids, cytochalasin B were obtained from Sigma (Poole,
10 Dorset, UK). Genistein for the sulphonylation reaction was obtained from Lancaster (Manchester UK).

**Synthesis of 2-Ethylestrone (2-EtE1)
and 2-Ethylestrone-3-*O*-sulfamate (2-EtEMATE)**

15

2-Ethylestrone (2-EtE1) and 2-Ethylestrone-3-*O*-sulfamate (2-EtEMATE) were prepared as follows see : briefly, Friedel-Crafts acetylation of estrone-3-*O*-methyl ether and catalytic hydrogenation followed by demethylation gave 2-Ethylestrone (2-EtE1) which was reacted with sulfonyl chloride to give the corresponding 3-*O*-sulfamate (2-EtEMATE). 2-
20 Ethylestradiol has been synthesised previously by a different route (21). Compounds were prepared as 10 mM stocks in tetrahydrofuran (THF).

2-Acetyestrone 3-methyl ether (8)

25 To a suspension of anhydrous aluminium chloride (3.76 g, 28.20 mmol) and acetyl chloride (2.0 ml, 28.13 mmol) and anhydrous nitromethane/dichloromethane (50 ml) at 0 °C, estrone methyl ether (4.0 g, 14.07 mmol) was added. After being stirred for 5 h at room temperature. The reaction mixture was poured into 10% HCl (100 ml) and the resulting mixture was extracted with ethyl acetate (3x100 ml). The combined organic extracts were
30 washed with brine until neutral, dried (MgSO₄), filtered and evaporated. The yellow/brown residue that was obtained was triturated with methanol and the resulting white precipitate was collected by filtration and air-dried to give a white solid (4.3 g) which was recrystallized from methanol to give 8 as white crystals (4.0 g, 87%); mp 181-184°C

(lit. 189-190°C);²²⁴ TLC (chloroform/acetone, 8:1): R_f 0.82; ν_{\max} (KBr) 1730 (C-17, C=O), 1670 (acetyl, C=O) cm^{-1} ; δ_{H} (400 MHz, CDCl_3) 0.9 (3H, s, C-18- CH_3), 1.39-2.54 (13H, m), 2.61 (3H, s, CH_3CO), 2.94 (2H, t, $J = 4.4$ Hz, C-6- H_2), 3.88 (3H, s, OCH_3), 6.69 (1H, s, C-4- H), and 7.7 (1H, s, C-1- H); δ_{C} (400 MHz, CDCl_3) 13.83 (q, C-18), 21.56 (t), 25.82 (t), 26.3 (q, CH_3CO), 29.9 (t), 31.44 (t), 31.93 (t), 35.84 (t), 38.24 (d), 43.75 (d), 47.95 (s, C13), 50.33 (d), 55.5 (q, OCH_3), 111.81 (d, C-4), 127.66 (d, C-1), 125.67 (s), 132.08 (s), 143.21 (s), 157.15 (s, C-3), 199.39 (s, C=O) and 220 (s, C-17-C=O). MS m/z (FAB+) 327.2 [100, (M+H)⁺], 311.2 (10), 173.2 (5). Found C, 77.4; H, 8.04 $\text{C}_{21}\text{H}_{26}\text{O}_3$ requires C, 77.27; H, 8.03%.

10

2-Ethylestrone 3-methyl ether (9)

A solution of 8 (1.0 g, 3.063 mmol) in THF/ethanol (1:2) (30 ml) was hydrogenated in the presence of Pd-C (1.0 g, 10%) 50 psi at room temperature for 24 h. After the supported catalyst was removed by filtration the filtrate was evaporated to give a white solid (950 mg) which was fractionated by flash chromatography (chloroform/acetone gradient). The less polar fraction gave a white solid (805 mg) which was further purified by recrystallization from methanol to give 9 as white crystals (775 mg, 78%); mp 112-115 °C; TLC (chloroform, and chloroform/acetone, 8:1): R_f s 0.56 and 0.78 respectively; ν_{\max} (KBr) 1730 (C=O), 1600 cm^{-1} ; δ_{H} ((400 MHz, CDCl_3), CDCl_3) 0.91 (3H, s, C-18- CH_3), 1.17 (3H, t, $J = 7.7$ Hz, CH_3CH_2), 1.39-2.48 (13H, m), 2.62 (2H, q, C-2- CH_2CH_3), 2.87 (2H, m, C-6- H_2), 3.81 (3H, s, OCH_3), 6.58 (1H, s, C-4- H), and 7.08 (1H, s, C-1- H); δ_{C} (400 MHz, CDCl_3) 13.86 (q), 14.54 (q), 21.58 (t), 22.86 (t), 25.82 (t), 26.17 (t), 29.72 (t), 31.63 (t), 35.88 (t), 38.23 (d), 44.04 (d), 48.23 (s, C13), 50.38, 55.32 (q, OCH_3), 110.8 (d, C-4), 126.34 (d, C-1), 130.08 (s), 131.31 (s), 134.71 (d), 155.41 (s) and 220 (C-17, C=O); MS m/z (FAB+) 312.1 [100, (M)⁺]. Found C, 80.5; H, 9.03 $\text{C}_{21}\text{H}_{28}\text{O}_2$ requires C, 80.73; H, 9.03%.

2-Ethylestrone (10)

30

Aluminum chloride (1.28 g, 9.615 mmol), sodium iodide (1.44 g, 9.615 mmol) and 9 (300 mg, 961.5 μmol) were added in this order to a mixture of acetonitrile (25 ml) and

dichloromethane (12.5 ml) at 0 °C under N₂. The resulting suspension was heated under reflux for 8 h, cooled to room temperature and then poured into water and followed by extraction with dichloromethane (3 x 100 ml). The combined DCM extracts were washed with 10% sodium thiosulphate (100 ml), brine, dried (MgSO₄), filtered and evaporated to give a yellow solid (290 mg), which was fractionated by flash chromatography (chloroform/acetone gradient). The yellow solid that isolated (254 mg) was further purified by recrystallization from acetone to give **10** as yellow crystals (240 mg, 83%); mp 201-204 °C; TLC (chloroform and chloroform/acetone, 8:1): R_f 0.3 and 0.65 respectively; ν_{max} (KBr) 3300 (OH), 1720 (C=O) cm⁻¹; δ_H (400 MHz, CDCl₃) 0.91 (3H, s, C-18-CH₃), 1.22 (3H, t, CH₃CH₂-), 1.38-2.51 (13H, m), 2.63 (2H, q, C-2-CH₂CH₃), 2.86 (2H, m, C-6-H₂), 4.68 (1H, br s, exchanged with D₂O, OH), 6.52 (1H, s, C-4-H) and (1H, s, C-1-H); MS *m/z* (FAB+) 298.0 [100, (M)⁺], 271.9 (10), 255.1 (10); MS *m/z* (FAB-) 297.1 [100, (M-H)⁻], 276.0 (45), 258.0 (40); Acc. mass (FAB+) 298.1928 requires C₂₀H₂₆O₂ 298.1933.

15 2-Ethylestrone 3-O-sulphamate (**11**)

To a stirred solution of **10** (150 mg, 502.6 μmol) and 2,6-di-*t*-butyl-4-methylpyridine (DBMP) (310 mg, 1.51 mmol) in CH₂Cl₂ (10 ml), sulphonoyl chloride in toluene (3.016 mmol) was added dropwise. After being stirred for 2 h the reaction mixture was diluted with dichloromethane (100 ml) and the resulting mixture washed with brine, dried (MgSO₄), filtered and evaporated. The residue that was obtained (170 mg) was fractionated by flash chromatography (chloroform/acetone gradient) and the white solid that isolated (155 mg) was further purified by recrystallization from ethyl acetate/hexane (1:2) to give **11** as white crystals (146 mg, 77%); mp 165-167 °C; TLC (chloroform/acetone, 8:1): R_f 0.41; ν_{max} (KBr) 3500-3300 (NH₂), 1720 (C=O), 1390 (SO₂N) cm⁻¹; δ_H (400 MHz, CDCl₃) 0.91 (3H, s, C-18-CH₃), 1.22 (3H, t, CH₃CH₂-), 1.41-2.55 (13H, m), 2.71 (2H, q, C-2-CH₂CH₃), 2.89 (2H, m, C-6-H₂), 5.0 (2H, br s, exchanged with D₂O, OSO₂NH₂), 7.11 (1H, s, C-4-H) and 7.2 (1H, s, C-1-H); MS *m/z* (FAB+) 531.2 [10, (M+H+NBA)⁺], 377.1 [100, (M)⁺], 298.2 [20, (M+H-SO₂NH₂)⁺]; MS *m/z* (FAB-) 530.1 [30, (M+NBA)⁻], 376.1 [100, (M-H)⁻], 275.1 (10); Acc. MS (FAB+) 377.1683 requires C₂₀H₂₇NO₄S 377.1661. Found C, 63.4; H, 3.73; N, 7.4 requires C₂₀H₂₇NO₄S C, 63.63; H, 3.71; N, 7.21%.

EXAMPLE 1 - Cell Culture

MCF-7 (ER+) and MDA-MB-231 (ER-) breast cancer cells were obtained from the American Type Culture Collection (Rockville, MD). Cells were routinely cultured in 25cm² culture flasks in Eagle's minimum essential medium (EMEM) with Hepes buffer (20mM). This medium was supplemented with L-glutamine (2mM), sodium hydrogen carbonate (10mM), 1% non-essential amino acids and 5% (v/v) foetal calf serum (FCS). Before adding test compounds, cells were washed with phosphate-buffered saline (PBS) and treatments added in phenol-red free medium containing 2% stripped FCS and supplements. The effects of 2-MeOE1 or 2-MeOEMATE on the growth of MCF-7 cells was assessed using a Cell Titer 96 cell proliferation assay (Promega, Southampton, Hants, UK) according to the manufacturers' instructions. For this, cells (5000 per well) were cultured in medium containing phenol-red and 10% FCS and were exposed to a drug for 4 days before the assay was performed. For MDA-MB-231 cells, cell numbers were determined using a Coulter counter.

For the culture of fibroblasts, resected breast tumour tissue was minced and incubated in EMEM for 18-24h at 37°C with collagenase (200µg/ml). The dispersed cells were harvested by centrifugation and washed twice with medium to remove collagenase. Dispersed cells were seeded into culture flasks and grown to confluence before passaging on a weekly basis. For experimental purposes 12 well multi-well plates or 25cm² flasks were seeded with fibroblasts and grown to 70-80% confluency. Cells were washed with PBS and exposed to drugs for 24h before determining cell numbers using a Coulter counter.

Photomicrographs of control and treated cells were taken under normal conditions of light and exposure using an Olympus SL35 Type 12 camera under an Olympus CK2 microscope (x100 magnification).

Results

The effect of 2-MeOE1 or 2-MeOEMATE on the proliferation of MCF-7 cells over a 4-day period was determined using a microwell plate proliferation assay (Fig 2). 2-MeOE1 at 0.1µM or 1.0µM had little effect on the proliferation of MCF-7 cells but reduced growth by 19% at 10µM. In contrast, 2-MeOEMATE inhibited cell proliferation by 29% and 52%

respectively at 0.1 μ M and 1.0 μ M. However, no further effect on cell proliferation was detected when cells were exposed to a higher (10 μ M) concentration of 2-MeOEMATE.

Exposure of MCF-7 cells to 2-MeOEMATE had a marked effect on cell morphology (Fig 3). In untreated cells (Fig 3a) only a few rounded cells were visible per field, whereas for cells treated with 2-MeOEMATE (1 μ M) for 24h there was a significant increase in the number of rounded and detached cells (Fig 3b). In contrast, 2-MeOE1 (1 μ M) had little effect on cell morphology (Fig 3c). Oestrone or EMATE when tested at 10 μ M did not affect cell morphology (data not shown).

2-MeOEMATE (1 μ M) had a similar effect on ER negative MDA-MB-231 breast cancer cell morphology and number (Figs 4a and 4b). At 1 μ M the number of MDA-MB-231 cells was reduced by 21% compared with untreated cells. As the stromal compartment constitutes a major proportion of the volume of breast tumours the effect of 2-MeOEMATE on the morphology of breast tumour-derived fibroblasts was also examined (Fig 5). At 1 μ M little effect on cell morphology was detected (Fig 5b) whereas at 5 μ M rounding of a significant proportion of fibroblasts occurred (Fig 5c). In contrast to the effect of 2-MeOEMATE on epithelial cell numbers it did not reduce the number of fibroblasts.

EXAMPLE 2 - Flow cytometry analysis

Cells were cultured in the presence of 2-MeOE1, 2-MeOEMATE or vehicle for up to 48h. To examine the reversibility of the effects of 2-MeOEMATE on the cell cycle, cells were treated for 24h with the drug after which cells were washed and cultured in fresh medium without drug for a further 24h period.

To prepare cells for analysis of DNA content, cells were trypsinised (0.25% trypsin, 0.02% EDTA), washed with PBS and fixed with 70% ethanol. Cells were collected by centrifugation, re-suspended in PBS (1-2 x 10⁶ cells/ml) and treated with RNase A (0.1mg/ml) and stained with propidium iodide (0.05mg/ml) for 30 min at room temperature. Cells were analysed with a flow cytometer (FACscan, Becton Dickinson Immunocytometry System, Bedford MA).

Results

MCF-7 cells were treated with 2-MeOEMATE (10 μ M) and the cell cycle distribution was analysed by flow cytometry. A time-course study revealed a progressive accumulation of cells in the G₂/M phase and the effect was apparent by 12h after treatment (Fig 6).

The proportion of cells in the G₁ phase of the cell cycle decreased from 66% for untreated cells to 50% and 23% respectively after 12h or 24h exposure to 2-MeOEMATE. There were corresponding increases in the proportions of cells in the G₂/M phase with little change in cells in the S phase being detected.

In a reversibility study, MCF-7 cells exposed to 2-MeOEMATE (10 μ M) for 24h or 48h again showed a marked increase in the proportion of cells in the G₂/M phase of the cell cycle (Fig 7a-d). For cells exposed to drug for 24h after which drug was removed and cells cultured in drug-free medium for a further 24h, a significant proportion of cells remained arrested in the G₂/M phase (Fig 7e). Quantitation of cells in the sub-G₁, G₁, S or G₂/M phase of the cycle (Table 1) confirmed the reciprocal decrease in the proportion of cells in the G₁ phase and increase in the G₂/M phase. However, a significant increase in the proportion of cells in the sub-G₁ fraction was also detected. Cells in the sub-G₁ fraction may represent cells undergoing apoptosis.

Table 1: Effect of 2-MeOEMATE (10 μ M) on cell cycle distribution

	Sub-G ₁	G ₁	S	G ₂ /M
Control 24h	9	51	21	19
2-MeOEMATE 24h	22	10	8	60
2-MeOEMATE 24h + 24h drug-free culture	27	8	7	57
Control 48h	4	76	8	12
2-MeOEMATE 48h	18	14	9	58

EXAMPLE 3 - TUNEL analysis

The ability of 2-MeOEMATE to induce apoptosis in MCF-7 cells was examined by TUNEL analysis using an *in situ* cell death detection kit (Boehringer Mannheim UK Ltd.,
5 Lewes, East Sussex, UK). Cells were fixed and permeabilised according to the manufacturers' instructions. Stained apoptotic cells were quantitated by flow cytometry.

Results

10 The possibility that cells in the sub-G₁ fraction may represent cells undergoing apoptosis was confirmed in a further experiment by TUNEL analysis (Fig 8). For untreated cells no increase in the proportion of fluorescently labelled cells was detected after staining. In contrast, there was a significant increase in the proportion of fluorescently labelled cells after exposure to 2-MeOEMATE (10µM) for 48h. Fluorescently labelled cells represented
15 approximately 10% of the cell population. This result indicates that 2-MeOEMATE can induce cells to undergo apoptosis.

EXAMPLE 4 - Effect of 2-MeOEMATE on growth of NMU-induced mammary tumours in intact rats

20

The effect of 2-MeOEMATE on mammary tumour growth was examined in a preliminary study using Ludwig rats (Harlan Olac, Bicester, UK) in which tumours were induced by inoculation of NMU. Tumour development was monitored regularly and when 0.5-1.0cm³ in volume, animals received vehicle (propylene glycol, 200µl/day, p.o.), 2-MeOEMATE
25 (20mg/kg/day, p.o.) or 2-MeOE1 (20mg/kg/day, p.o.) daily for an 11 day period. Tumour length and width was measured with callipers and tumour volumes calculated as described (21).

Results

30

A preliminary study was carried out to compare the abilities of 2-MeOE1 and 2-MeOEMATE to inhibit tumour growth *in vivo*. For this, the growth of mammary tumours was initiated by inoculation with NMU. Drugs were administered orally when tumour

volumes reached 0.5-1.0cm³. For two of the animals receiving vehicle, tumour volumes continued to increase (average 82%) while little change in the volume of a tumour in a third animal was detected (Fig 9). For two animals receiving 2-MeOE1 no change in tumour volume occurred in one, while for the other a modest (25%) reduction was detected
5 over the 11-day period of the study.

For three animals receiving 2-MeOEMATE the tumour volume in one animal continued to increase up to day 6, but thereafter showed a slight (8%) reduction. In contrast, for the two other animals receiving 2-MeOEMATE, tumours regressed rapidly and were barely
10 palpable at the end of the 11-day period. Tumour volumes in the two animals receiving 2-MeOEMATE that regressed were monitored for a further 33 days during which time no regrowth of tumours was detected.

EXAMPLE 5 - Irreversible Effect of 2-Methoxy- or 2-Ethylloestrone Sulphamates on 15 Growth of MCF-7 Breast Cancer Cells

Procedure

Stage 1

20 MCF-7 breast cancer cells were seeded into multi-well culture plates at a density of 10⁵ cells/well. Cells were allowed to attach and grown until about 30% confluent when they were treated as follows:

Control - no treatment

25

2-MeOE1 5µm

2-MeOE1 1µm

2-MeOEMATE 5µm

30

2-MeOEMATE 1µm

2-EtE1 5µm

2-EtE1 1µm

2-EtEMATE 5 μ m

2-EtEMATE 1 μ m

- 5 Cells were grown for 6 days in growth medium containing the above drugs with changes of medium/drug every 3 days. At the end of this period cell numbers were counted using a Coulter cell counter.

The results from this first stage of the experiment are shown in Fig 10. Compared with the
10 controls, 2-methoxyoestrone or 2-ethyloestrone had little effect on cell growth. In contrast, treatment of cells with 2-methoxyoestrone sulphamate at 5 μ m or 1m reduced cell numbers to 43% and 52% of the control cell number respectively. The corresponding values after treatment of cells with 2-ethyloestrone sulphamate were 36% and 50% respectively.

15 Stage 2

After treatment of cells for a 6-day period with the above drugs cells were re-seeded at a density of 10⁴ Cells/well. No further treatments were added. Cells were allowed to
20 continue to grow for a further 6 days in the presence of growth medium. At the end of this period cell numbers were again counted.

Results from this part of the experiment are shown in Fig 11. Compared with the controls 2-methoxyoestrone or 2-ethyloestrone had little effect on cell numbers showing that these
25 compounds have no irreversible growth inhibitory effects on these cells.

In contrast, cells treated with either 2-methoxyoestrone sulphamate or 2-ethyloestrone sulphamate were severely growth restricted. These results demonstrated that once cells
30 have been exposed to 2-methoxyoestrone sulphamate or 2-ethyloestrone sulphamate their growth is irreversibly compromised.

EXAMPLE 6 - Effect of Non-Hydrocarbyl/Oxyhydrocarbyl Substituted Sulphamates on Growth of MCF-7 Breast Cancer Cells

Procedure

Stages 1 and 2 described above were repeated using Control - no treatment, EMATE 20 μ M and EMATE 5 μ M.

5 The stage 1 and 2 results were:

Stage 1 EMATE 20 μ M = 119% control

EMATE 5 μ M = 139% control

Stage 2 EMATE 20 μ M = 103% control

10 EMATE 5 μ M = 98% control

These data show the importance of the hydrocarbyl/oxyhydrocarbyl substituents on the compounds of the present invention to achieve inhibition and/or prevention and/or arrest of cell cycling.

15

EXAMPLE 7 - Cell cycle and Apoptosis Analysis

MCF7, CAL51, CAMA1 and ZR-75-1 breast cancer derived cell lines were obtained from ATCC (MCF7, CAMA1, ZR-75-1) or from the Dutrillaux laboratory (CAL51) (22) and
20 maintained in Dulb cco's modified Eagle's medium (DMEM) supplemented with 10% (v/v) fetal calf serum and antibiotics.

DNA content was determined by flow cytometric analysis of propidium iodide stained cells and TdT-mediated dUTP-nick end labelling (TUNEL), respectively, as described (20). The
25 proportion of cells in G2/M phase of the cell cycle was calculated as a proportion of cells with 2N to 4N DNA content. The proportion of cells with <G1 DNA content was calculated as a percentage of total cells.

To determine the proportion of cells in mitosis, drug treated cells were collected by
30 trypsinisation and cytopspins prepared. Cells were fixed in ice cold methanol for 5 minutes, air dried and DNA was stained using propidium iodide (PI) (0.1 mg/ml in phosphate buffered saline (PBS) containing 10% (v/v) newborn calf serum and 0.05% (w/v) sodium

azide). Cells were analysed by confocal microscopy using a Zeiss Axiovert 100 M microscope equipped with the LSM 510 confocal system (Zeiss, Jena, Germany).

We examined whether the differential sensitivity of breast cancer cell lines to the growth inhibitory effects of sulfamoylated estrones was related to differences in extent/phase of cell cycle arrest and/or cell death. These studies focused on 2-EtEMATE since we were readily able to synthesise relatively large amounts of this compound. We first examined the effects of 2-EtEMATE on the DNA content of MCF7, ZR-75-1, CAL51 and CAMA1 cells using flow cytometry of propidium iodide (PI) stained cells (Fig. 13 and Table 2). Cells were treated with 2-EtEMATE at 500 nM since this was the lowest dose that gave significant growth inhibition in each cell line (Fig. 14). MCF7 cells accumulated in the G2/M phase of the cell cycle within 24 hours and cells were maximally arrested (approximately 60% of cells) after 48 hours. This arrest was maintained for the duration of the experiment although there was a modest increase in cells with <G1 DNA content indicative of cell death at 72 and 96 hours. Similar results were obtained in ZR-75-1 cells although G2/M arrest was delayed in these cells (60% cells in G2/M after 72 hours). By contrast, CAL51 and CAMA1 cells which were more sensitive than MCF7 or ZR-75-1 cells in the microtitre plate assay (Fig. 14) underwent a more rapid G2/M arrest (approximately 60% of cells were in G2/M within 24 hours). In addition, 2-EtEMATE induced significant cell death within 48 hours and approximately half of the cells were dead after 96 hours (Table 2).

Table 2. Effect of 2-EtEMATE on DNA content of breast cancer cell lines. Cell cycle parameters of MCF7, ZR-75-1, CAL51 and CAMA1 cells exposed to 2-EtEMATE (500 nM) for up to 96 hours. The proportion of cells in G1/S or G2/M are shown as a percentage of total cells with a 2N to 4N DNA content. The proportion of cells with a sub-G1 DNA content is shown as percent of total cells. Untreated cells are at 96 hours.

Cell Line	MCF7			ZR-75-1		
	<G1	G1/S	G2/M	<G1	G1/S	G2/M
Untreated	2	77	23	1	79	20
24 hours	5	63	37	1	84	15
48 hours	5	38	62	6	60	40
72 hours	12	34	66	12	37	63
96 hours	8	29	71	9	20	80

Cell Line	CAL51			CAMA1		
	<G1	G1/S	G2/M	<G1	G1/S	G2/M
Untreated	1	78	22	4	70	30
24 hours	2	38	62	4	33	67
48 hours	10	18	82	16	25	75
72 hours	21	32	68	36	33	67
96 hours	44	23	77	51	33	67

Flow cytometric analysis demonstrated that cells treated with the sulfamoylated estrone derivatives were arrested in G2/M phases of the cell cycle. Cells treated with 2-EtEMATE or 2-MeOEMATE often had a characteristic "rounded-up" morphology (Ref. 20 and data not shown) suggesting that cells were in fact arrested in mitosis. To determine whether cells were in interphase or mitosis, we stained drug-treated MCF7 cells with PI to visualise chromosomes. In preliminary experiments, we found that the "rounded up" cells did not fix to the collagen-treated glass slides that we used for fluorescence staining experiments and we therefore collected all of the cells by trypsinisation and prepared cytopspins prior to staining. The vast majority of control MCF7 cells had a uniformly stained nucleus characteristic of interphase cells with uncondensed chromosomes (Fig. 15 and Table 3). By contrast, a significant proportion of cells treated with 2-EtEMATE or 2-MeOEMATE showed condensed chromosomes characteristic of mitosis. Therefore, the sulfamoylated estrone derivatives induce a mitotic arrest. Consistent with lack of effect in FACs assay (Fig. 13), the non-sulfamoylated estrone derivatives did not increase the number of mitotic cells. Since it was necessary to use cytopspins in these experiments, it was difficult to determine the architecture of the chromosomes in cells treated with sulfamoylated estrone derivatives. However, the chromosomes appeared to be fully condensed suggesting that cells had reached pro-metaphase/metaphase.

Table 3. Effect of estrone derivatives on mitosis in MCF7 cells. MCF7 cells were treated with the indicated compounds for 24 hours. Cells were recovered by trypsinisation and cytopspins prepared. DNA was stained with PI and cells in mitosis (i.e., with condensed chromosomes) determined as a percentage of total cells.

Drug	mitotic cells
Control	4%
2-EtEMATE, 5 μ M	49%

	2-EtEMATE, 500 nM	29%
	2-MeOEMATE, 5 μ M	38%
	2-MeOEMATE, 500 nM	26%
	2-EtE1, 5 μ M	6%
5	2-EtE1, 500 nM	4%

We used the TUNEL assay to confirm that the cell death detected by flow cytometry of PI stained cells was due to apoptosis. Following exposure to 2-EtEMATE (500 nM for 72 hours) there was a significant increase in the proportion of CAL51 and CAMA1 cells
10 undergoing apoptosis (60% and 37% TUNEL positive cells, respectively) (Fig. 16). By contrast, there was only a modest increase in TUNEL positivity in MCF7 or ZR-75-1 cells (7% and 10%, respectively) treated with 2-EtEMATE at this concentration.

Taken together, these analyses demonstrate that the variations in the sensitivity of breast
15 cancer cell lines to short term growth inhibition by sulfamoylated estrones are reflected in differences in effects on cell cycle and apoptosis. Although 2-EtEMATE induced a G2/M arrest in all cell lines, this was more rapid in relatively sensitive ER negative CAL51 and CAMA1 cells than in ER positive MCF7 and ZR-75-1 cells. Furthermore, 2-EtEMATE was a more potent inducer of apoptosis in CAL51 and CAMA1 cells than MCF7 and ZR-
20 75-1 cells.

DISCUSSION

Our results add further weight to the evidence showing that 2-methoxyoestrogens, or their
25 synthetic analogues (23, 24), represent a new class of drugs for cancer therapy. The present compounds, such as 2-MeOEMATE, as previously found for 2-MeOE2, had a marked effect on the morphology and growth of MCF-7 and MDA-MD-231 breast cancer cells. Fibroblasts derived from breast tumours also showed a similar rounding in response to 2-MeOEMATE although a higher concentration was required to induce morphological
30 changes in these cells.

The marked effect that 2-MeOEMATE has on the growth of ER+ and ER- breast cancer cells confirms that the present compounds such as 2-methoxyoestrogen sulphamates should

be active against both hormone-dependent and independent breast tumours. At 1 μ M 2-MeOE1 had little effect on the proliferation of MCF-7 cells while 2-MeOEMATE, at this concentration, inhibited proliferation by 52%. The reason for the increased potency conferred by the addition of a sulphamate group to 2-MeOE1 in the *in vitro* assays is not readily apparent. From *in vivo* studies with oestrogen sulphamates, it is known that these compounds are released slowly from rbc's to give a protracted increase in steroid blood concentration (17). Other *in vivo* studies have indicated that EMATE is capable of inactivating steroid sulphatase for a prolonged period of time after a single dose or multiple doses (25). Thus, it is likely that EMATE is binding to a cellular protein from which it is slowly released. This may account for the enhanced potency of the sulphamate *in vitro* compared with that of 2-MeOE1 in being able to reduce cell proliferation.

From the DNA analysis it is apparent that the present compounds, e.g. 2-MeOEMATE, like 2-MeOE2 and taxol, induce an arrest of cell cycling, in particular an arrest of cells in the G₂/M phase of the cell cycle (9, 26, 27). For 2-MeOE2, however, a washout experiment, in which cells continued to be cultured in drug-free medium after an initial period of exposure to the drug, revealed that a significant proportion of cells re-entered the G₁/S phase of the cell cycle by 24h after removal of the drug (13). In contrast, for the compounds of the present invention (2-MeOEMATE) cells remained arrested in the G₂/M phase for at least 24h after removal of the drug. This finding provides further evidence that the present compounds may be binding to a cell protein.

2-MeOEMATE, while possessing novel anti-proliferative effects, remains a potent steroid sulphatase inhibitor (18).

Similar experiments in which MCF-7 cells were exposed to EMATE and then extensively washed to remove drug revealed that steroid sulphatase remained almost completely inactivated (28).

The ability of 2-MeOE2 to inhibit the growth of MCF-7 cells and to induce cells to become rounded and detached has previously been shown to result from its ability to induce apoptosis in these cells (8, 9). In the present investigation a significant increase in cells in the sub-G₁ fraction was detected after exposure to a present compound, namely 2-

MeOEMATE. Cells in this fraction are thought to represent cells that have undergone apoptosis. TUNEL analysis confirmed that 2-MeOEMATE did induce a proportion of cells to undergo apoptosis. Like 2-MeOE2 and other drugs, such as taxol, that alter microtubule stability, the present compounds (2-MeOEMATE) probably induces apoptosis by increasing the phosphorylation of the oncoprotein Bcl-2 (27-29). Bcl-2 belongs to a family of proteins that are anti-apoptotic and their ability to inhibit apoptosis from their dimerisation with, and inactivation of proapoptotic proteins such as Bax (30, 31). Phosphorylation of Bcl-2 blocks its ability to dimerise with Bax thus allowing the induction of apoptosis. It has been clearly demonstrated that phosphorylation of Bcl-2 occurs during the arrest of cells in the G₂/M phase of the cell cycle (30). In a preliminary study 2-MeOE2 was found to induce phosphorylation of Bcl-2 in leukaemia cells (29). This indicates that 2-methoxyoestrogens have a similar mechanism of action to that of taxol and other drugs that cause microtubule damage.

The marked effect that a present compound, especially 2-MeOEMATE, had on cell proliferation *in vitro* led to a preliminary *in vivo* study. *In vivo* an NMU-induced mammary tumour in one of the animals receiving 2-MeOE1 showed a modest (25%) regression. In contrast, 2/3 tumours in animals receiving 2-MeOEMATE regressed almost completely by the end of the 11-day study. The enhanced efficacy of 2-MeOEMATE compared with that of 2-MeOE1 lends support to the findings from the *in vitro* studies that the sulphamoylated estrogen is more potent than its parent compound.

Other *in vivo* tumour studies with the present compounds (2-methoxyoestrogens) have also produced encouraging results although at much higher than the doses employed in the present study. 2-MeOE2 (100mg/kg, p.o.) when administered every other day to C₅₇BL/6 mice significantly reduced the growth of subcutaneously inoculated Meth A sarcoma and B16 melanoma tumours by 57% and 83% respectively (10). As in our study, this response was achieved in a relatively short, 12 day, period of time. In addition to inhibiting the growth of these tumours, tumour neovascularisation was markedly reduced. This suggests that an important action of 2-methoxyoestrogens is inhibiting tumour growth. Oral administration of 2-MeOE2 (75mg/kg/day for 29 days) also suppressed the growth of tumours resulting from inoculation of the ER- MDA-MB-435 breast cancer cells in SCID mice (11). Tumour volumes were reduced by 60% compared with those of untreated controls. No toxic side

effects were detected at this relatively high dosage of the drug. In addition to these animal studies, 2-MeOE2 is currently undergoing a Phase I/II trial although details of the outcome of this trial are not yet available (42, 43).

- 5 In the present *in vivo* studies, 2-MeOEMATE was employed although *in vitro* investigation has found that 2-MeOE2 is more potent than 2-MeOE1 (10). However, from experiments with estradiol sulphamate it is known that the 17-hydroxyl function is oxidised during gastric transit yet the reduced form of the steroid is released from rbc's (17). Therefore, there would appear to be no advantage to be gained from the oral administration of the
10 reduced form of 2-MeOEMATE.

As most cancers eventually become resistant to either hormone or chemotherapy the development of drugs that act on different cellular targets offers considerable hope for the development of new cancer therapies. The therapeutic use of taxol has been an important
15 advance but problems with its solubility and toxicity have limited its prolonged use. The identification of 2-MeOEMATE as a form of 2-methoxyoestrogen with enhanced efficacy, bioavailability and duration of action suggests that this drug should have considerable potential for cancer therapy.

- 20 Thus, in summary, the present invention provides a composition and compound suitable for use in the treatment of cancers and, especially, breast cancer.

In particular, in one aspect the present invention addresses the problem of blocking the growth of cancers including leukaemias and solid tumours such as breast, endometrium,
25 prostate, ovary and pancreatic tumours.

It is also believed that the present invention has implications in treating hormonal conditions in addition to those associated with oestrogen. Hence, the present invention also provides a composition that is capable of affecting hormonal activity and is capable of
30 affecting an immune response, wherein the composition is the composition of the present invention.

It is also to be understood that the compound/composition of the present invention may

have other important medical implications.

For example, the composition of the present invention may be useful in the treatment of the disorders listed in WO-A-98/05635. For ease of reference, part of that list is now
5 provided: cancer, inflammation or inflammatory disease, dermatological disorders, fever, cardiovascular effects, haemorrhage, coagulation and acute phase response, cachexia, anorexia, acute infection, HIV infection, shock states, graft-versus-host reactions, autoimmune disease, reperfusion injury, meningitis, migraine and aspirin-dependent anti-thrombosis; tumour growth, invasion and spread, angiogenesis, metastases, malignant,
10 ascites and malignant pleural effusion; cerebral ischaemia, ischaemic heart disease, osteoarthritis, rheumatoid arthritis, osteoporosis, asthma, multiple sclerosis, neurodegeneration, Alzheimer's disease, atherosclerosis, stroke, vasculitis, Crohn's disease and ulcerative colitis; periodontitis, gingivitis; psoriasis, atopic dermatitis, chronic ulcers, epidermolysis bullosa; corneal ulceration, retinopathy and surgical wound healing; rhinitis,
15 allergic conjunctivitis, eczema, anaphylaxis; restenosis, congestive heart failure, endometriosis, atherosclerosis or endosclerosis.

In addition, or in the alternative, the composition of the present invention may be useful in the treatment of disorders listed in WO-A-98/07859. For ease of reference, part of that list
20 is now provided: cytokine and cell proliferation/differentiation activity; immunosuppressant or immunostimulant activity (e.g. for treating immune deficiency, including infection with human immune deficiency virus; regulation of lymphocyte growth; treating cancer and many autoimmune diseases, and to prevent transplant rejection or induce tumour immunity); regulation of haematopoiesis, e.g. treatment of myeloid or
25 lymphoid diseases; promoting growth of bone, cartilage, tendon, ligament and nerve tissue, e.g. for healing wounds, treatment of burns, ulcers and periodontal disease and neurodegeneration; inhibition or activation of follicle-stimulating hormone (modulation of fertility); chemotactic/chemokinetic activity (e.g. for mobilising specific cell types to sites of injury or infection); haemostatic and thrombolytic activity (e.g. for treating haemophilia and stroke);
30 antiinflammatory activity (for treating e.g. septic shock or Crohn's disease); as antimicrobials; modulators of e.g. metabolism or behaviour; as analgesics; treating specific deficiency disorders; in treatment of e.g. psoriasis, in human or veterinary medicine.

In addition, or in the alternative, the composition of the present invention may be useful in the treatment of disorders listed in WO-A-98/09985. For ease of reference, part of that list is now provided: macrophage inhibitory and/or T cell inhibitory activity and thus, anti-inflammatory activity; anti-immune activity, i.e. inhibitory effects against a cellular and/or humoral immune response, including a response not associated with inflammation; inhibit the ability of macrophages and T cells to adhere to extracellular matrix components and fibronectin, as well as up-regulated fas receptor expression in T cells; inhibit unwanted immune reaction and inflammation including arthritis, including rheumatoid arthritis, inflammation associated with hypersensitivity, allergic reactions, asthma, systemic lupus erythematosus, collagen diseases and other autoimmune diseases, inflammation associated with atherosclerosis, arteriosclerosis, atherosclerotic heart disease, reperfusion injury, cardiac arrest, myocardial infarction, vascular inflammatory disorders, respiratory distress syndrome or other cardiopulmonary diseases, inflammation associated with peptic ulcer, ulcerative colitis and other diseases of the gastrointestinal tract, hepatic fibrosis, liver cirrhosis or other hepatic diseases, thyroiditis or other glandular diseases, glomerulonephritis or other renal and urologic diseases, otitis or other oto-rhino-laryngological diseases, dermatitis or other dermal diseases, periodontal diseases or other dental diseases, orchitis or epididymo-orchitis, infertility, orchidial trauma or other immune-related testicular diseases, placental dysfunction, placental insufficiency, habitual abortion, eclampsia, pre-eclampsia and other immune and/or inflammatory-related gynaecological diseases, posterior uveitis, intermediate uveitis, anterior uveitis, conjunctivitis, chorioretinitis, uveoretinitis, optic neuritis, intraocular inflammation, e.g. retinitis or cystoid macular oedema, sympathetic ophthalmia, scleritis, retinitis pigmentosa, immune and inflammatory components of degenerative fundus disease, inflammatory components of ocular trauma, ocular inflammation caused by infection, proliferative vitreo-retinopathies, acute ischaemic optic neuropathy, excessive scarring, e.g. following glaucoma filtration operation, immune and/or inflammation reaction against ocular implants and other immune and inflammatory-related ophthalmic diseases, inflammation associated with autoimmune diseases or conditions or disorders where, both in the central nervous system (CNS) or in any other organ, immune and/or inflammation suppression would be beneficial, Parkinson's disease, complication and/or side effects from treatment of Parkinson's disease, AIDS-related dementia complex HIV-related encephalopathy, Devic's disease, Sydenham chorea, Alzheimer's disease and other degenerative diseases,

conditions or disorders of the CNS, inflammatory components of stokes, post-polio syndrome, immune and inflammatory components of psychiatric disorders, myelitis, encephalitis, subacute sclerosing pan-encephalitis, encephalomyelitis, acute neuropathy, subacute neuropathy, chronic neuropathy, Guillain-Barre syndrome, Sydenham chora, myasthenia gravis, pseudo-tumour cerebri, Down's Syndrome, Huntington's disease, amyotrophic lateral sclerosis, inflammatory components of CNS compression or CNS trauma or infections of the CNS, inflammatory components of muscular atrophies and dystrophies, and immune and inflammatory related diseases, conditions or disorders of the central and peripheral nervous systems, post-traumatic inflammation, septic shock, infectious diseases, inflammatory complications or side effects of surgery, bone marrow transplantation or other transplantation complications and/or side effects, inflammatory and/or immune complications and side effects of gene therapy, e.g. due to infection with a viral carrier, or inflammation associated with AIDS, to suppress or inhibit a humoral and/or cellular immune response, to treat or ameliorate monocyte or leukocyte proliferative diseases, e.g. leukaemia, by reducing the amount of monocytes or lymphocytes, for the prevention and/or treatment of graft rejection in cases of transplantation of natural or artificial cells, tissue and organs such as cornea, bone marrow, organs, lenses, pacemakers, natural or artificial skin tissue.

PROTOCOL I

IRREVERSIBLE INHIBITION

Procedure

Stage 1

MCF-7 breast cancer cells were seeded into multi-well culture plates at a density of 10^5 cells/well. Cells were allowed to attach and grown until about 30% confluent when they were treated as follows:

Control - no treatment

Compound of Interest (COI) $20\mu\text{M}$

Cells were grown for 6 days in growth medium containing the COI with changes of

medium/COI every 3 days. At the end of this period cell numbers were counted using a Coulter cell counter.

Stage 2

5

After treatment of cells for a 6-day period with the COI cells were re-seeded at a density of 10^4 cells/well. No further treatments were added. Cells were allowed to continue to grow for a further 6 days in the presence of growth medium. At the end of this period cell numbers were again counted.

10

All publications and patents mentioned in the above specification are herein incorporated by reference.

15

Various modifications and variations of the present invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific
20 embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in chemistry, biology or related fields are intended to be within the scope of the following claims.

References

1. Nebert, D.W. Proposed role of drug-metabolising enzymes: regulation of steady-state levels of the ligands that effect growth, homeostasis, differentiation and neuroendocrine function. *Molec. Endocrinol.*, 5: 1203-1214, 1991.
2. Reed, M.J., Purohit, A., Woo, L.W.L., and Potter, B.V.L. The development of steroid sulphatase inhibitors. *Endocr.-Rel. Cancer*, 3: 9-23, 1996.
3. Bradlow, H.L., Michnovicz, J.J., Telang, N.T., and Osborne, M.P. Effect of dietary indole-3-carbinol on oestradiol metabolism and spontaneous tumours in mice. *Carcinogenesis*, 12: 1571-1574, 1991.
4. Bradlow, H.L., Sepkovic, D.W., Telang, N.T., and Osborne, M.P. Indole-3-carbinol: a novel approach to breast cancer prevention. *Ann. N.Y. Acad. Sci.*, 728: 180-200, 1995.
5. Bradlow, H.L., Davis, D.L., Lin, G., Sepkovic, D.W., and Tiwari, R. Effects of pesticides on the ratio of 16 α /2-hydroxyoestrone: a biological marker of breast cancer risk. *Environ. Health Perspect.*, 103 (Suppl.): 147-150, 1995.
6. Bradlow, H.L., Telang, N.T., Sepkovic, D.W., and Osborne, M.P. 2-Hydroxyoestrone: the 'good' estrogen. *J. Endocrinol.*, 150: S259-S265, 1996.
7. Zhu, B.T., and Conney, A.H. Is 2-methoxyoestradiol an endogenous estrogen metabolite that inhibits mammary carcinogenesis. *Cancer Res.*, 58: 2269-2277, 1998.
8. Seegers, J.C., Aveling, M.-L., Van Aswegen, C.H., Cross, M., Koch, F., and Joubert, W.S. The cytotoxic effects of oestradiol-17 β , catecholoeestradiols and methoxyoestradiols on dividing MCF-7 and HeLa cells. *J. Steroid Biochem.*, 32: 797-809, 1989.
9. Seegers, J.C., Lottering, M.L., Grobler, C.J., Van Papendorp, D.H., Habbersett, R.C., Shou, Y., and Lehnert, B.E. The mammalian metabolite, 2-methoxyoestradiol affects p53 levels and apoptosis induction in transformed cells but not in normal cells. *J. Steroid Biochem. Molec. Biol.*, 62: 253-267, 1997.
10. Fotsis, T., Zhang, Y., Pepper, M.S., Adlercreutz, H., Montesano, R., Nawroth, P.P., and Schweigerer, L. The endogenous oestrogen metabolite 2-methoxyoestradiol inhibits angiogenesis and suppresses tumour growth. *Nature*, 368: 237-239, 1994.
11. Klauber, N., Parangi, S., Flynn, E., Hamel, E., and D'Amato, R.J. Inhibition of angiogenesis and breast cancer in mice by the microtubule inhibitors 2-methoxyoestradiol and taxol. *Cancer Res.*, 57: 81-86, 1997.
12. D'Amato, R.J., Lin, C.M., Flynn, E., Folkman, J., and Hamel, E. 2-

Methoxyoestradiol, an endogenous mammalian metabolite, inhibits tubulin polymerization by interacting at the colchicine site. *Proc. Natl. Acad. Sci. USA*, 91: 3964-3968, 1994.
AND US Patent No. 5,504,074

13. Attalla, H., Mäkelä, T.P., Adlercreutz, H., and Andersson, L.C. 2-Methoxyoestradiol
5 arrests cells in mitosis without depolymerising tubulin. *Biochem. Biophys. Res. Comm.*,
228: 467-473, 1996.
14. Howarth, N.M., Purohit, A., Reed, M.J., and Potter, B.V.L. Estrone sulphamates:
potent inhibitors of oestrone sulphatase with therapeutic potential. *J. Med. Chem.*, 37:
219-221, 1994.
- 10 15. Purohit, A., Williams, G.J., Howarth, N.M., Potter, B.V.L., and Reed, M.J.
Inactivation of steroid sulphatase by an active site-directed inhibitor, oestrone-3-*O*-
sulphamate. *Biochemistry*, 34 : 11508-11514, 1995.
16. Elger, W., Schwarz, S., Hedden, A., Reddersen, G., and Schneider, B. Sulphamates of
various oestrogens as prodrugs with increased systemic and reduced hepatic oestrogenicity
15 at oral application. *J. Steroid Biochem. Molec. Biol.*, 55: 395-403, 1995.
17. Elger, W., Palme, H.-J., and Schwarz, S. Novel oestrogen sulphamates: a new
approach to oral hormone therapy. *Exp. Opin. Invest. Drugs*, 7: 575-589, 1998.
18. Purohit, A., Vernon, K.A., Wagenaar-Humelinck, A.E., Woo, L.W.L., Hejaz, H.A.M.,
Potter, B.V.L., and Reed, M.J. The development of A-ring modified analogues of
20 oestrone-3-*O*-sulphamate as steroid sulphatase inhibitors with reduced oestrogenicity. *J.*
Steroid Biochem. Molec. Biol., 64: 269-275, 1998.
19. Purohit, A., Hejaz, H.A.M., Woo, L.W.L., Van Strien, A.E., Potter, B.V.L., and Reed,
M.J. Recent advances in the development of steroid sulphatase inhibitors. *J. Steroid*
Biochem. Molec. Biol. (In Press).
- 25 20. Woo, L.W.L., Purohit, A., Reed, M.J., and Potter, B.V.L. Active site directed
inhibition of oestrone sulphatase by non-steroidal coumarin sulphamates. *J. Med. Chem.*,
39: 1349-1351, 1996.
21. Wilkinson, J.R., Williams, J.C., Singh, D., Goss, P.E., Easton, D., and Coombes, R.C.
Response of nitrosomethylurea-induced rat mammary tumor to endocrine therapy and
30 comparison with clinical response. *Cancer Res.*, 46 : 4862-4865, 1986.
22. Vera, J.C., Reyes, A.M., Carcamo, J.G., Velasquez, F.V., Rivas, C.I., Zhang, R.H.,
Strobel, P., Iribarren, R., Scher, H.I., Slebe, J.C., and Golde, D.W. Genistein is a natural
inhibitor of hexose and dehydroascorbic acid transport through the glucose transporter,


- GLUT1. *J. Biol. Chem.*, 271 : 8719-8724, 1996.
23. Cushman, M., He, H.-M., Katzenellenbogen, J.A., Lin, C.M., and Hamel, E. Synthesis, antitubulin and antimitotic activity and cytotoxicity of analogues of 2-methoxyoestradiol, an endogenous mammalian metabolite of oestradiol that inhibits tubulin polymerisation by binding to the colchicine binding site. *J. Med. Chem.*, 38: 2041-2049, 1995.
24. Cushman, M., He, H.-M., Katzenellenbogen, J.A., Varma, R.K., Hamel, E., Lin, C.M., Ram, S., and Sachdeva, Y.P. Synthesis of analogues of 2-methoxyoestradiol with enhanced inhibitory effects on tubulin polymerisation and cancer cell growth. *J. Med. Chem.*, 40: 2323-2334, 1997.
25. Purohit, A., Williams, G.J., Roberts, C.J., Potter, B.V.L., and Reed, M.J. *In vivo* inhibition of oestrone sulphatase and dehydroepiandrosterone sulphatase by oestrone-3-O-sulphamate. *Int. J. Cancer*, 63: 106-111, 1995.
26. Haldar, S., Jena, N., and Croce, C.M. Inactivation of bcl-2 by phosphorylation. *Proc. Natl. Acad. Sci. USA*, 92: 4507-4511, 1995.
27. Haldar, S., Chintapalli, J., and Croce, C.M. Taxol induces bcl-2 phosphorylation and death of prostate cancer cells. *Cancer Res.*, 56: 1253-1255, 1996.
28. Purohit, A., Reed, M.J., Morris, N.C., Williams, G.J., and Potter, B.V.L. Regulation and inhibition of steroid sulphatase activity in breast cancer. *Ann. N.Y. Acad. Sci.*, 784: 40-49, 1996.
29. Attalla, H., Westberg, J.A., Andersson, L.C., Adlercreutz, H., and Makela, T.P. 2-Methoxyestradiol-induced phosphorylation of Bcl-2: uncoupling from JNK/SAPK activation. *Biochem. Biophys. Res. Comm.*, 247: 616-619, 1998.
30. Haldar, S., Basu, A., and Croce, C.M. Bcl-2 is the guardian of microtubule integrity. *Cancer Res.*, 57: 229-233, 1997.
31. Blagosklonny, M.V., Giannakakou, P., El-Deiry, W.S., Kingston, D.G.T., Higgs, P.I., Neckers, L., and Fojo, T. Raf-1/bcl-2 phosphorylation: a step from microtubule damage to cell death. *Cancer Res.*, 57: 130-135, 1997.
32. Constantinou, A.I., Kamath, N., and Murley, J.S. Genistein inactivates bcl-2, delays the G₂/M phase of the cell cycle and induces apoptosis of human breast adenocarcinoma MCF-7 cells. *Eur. J. Cancer*, 34: 1927-1934, 1998.
33. Akiyama, T., Ishida, J., Nakagawa, S., Ogawara, H., Watanabe, S.-I., Itoh, N., Shibuya, M., and Fukami, Y. Genistein, a specific inhibitor of tyrosine-specific protein kinases. *J.*

Biol. Chem., 262: 5592-5595, 1987.

34. Flier, J.S., Mueckler, M.M., Usher, P., and Lodish, H.F. Elevated levels of glucose transport and transporter messenger RNA are induced by *ras* or *src* oncogenes. *Science*, 235: 1492-1495, 1987.
- 5 35. Shim, H., Chun, Y.-S., Lewis, B.C., and Dang, C.V. A unique glucose-dependent apoptotic pathway induced by *c-myc*. *Proc. Natl. Acad. Sci. USA*, 95: 1511-1516, 1998.
36. Gould, G.W., and Holman, G.D. The glucose transporter family: structure, function and tissue-specific expression. *Biochem. J.*, 295: 329-341, 1993.
37. Brown, R.S., and Wahl, R.L. Overexpression of GLUT-1 glucose transporter in
10 human breast cancer. *Cancer*, 72: 2979-2985, 1993.
38. Younes, M., Lechago, L.V., Somoano, J.R., Mosharaf, M., and Lechago, J. Wide expression of the human erythrocyte glucose transporter GLUT-1 in human cancers. *Cancer Res.*, 56: 1164-1167, 1996.
39. Zamora-Leon, P., Golde, D.W., Concha, I.I., Rivas, C.I., Delgado-Lopez, F., Baselga,
15 J., Nualart, F., and Vera, J.C. Expression of the fructose transporter GLUT-5 in human breast cancer. *Proc. Natl. Acad. Sci.*, 93: 1847-1852, 1996.
40. Tsakiridis, T., Vranic, M., and Klip, A. Disassembly of the actin network inhibits insulin-dependent stimulation of glucose transport and prevents recruitment of glucose transporters to the plasma membrane. *J. Biol. Chem.*, 269: 29934-29942, 1994.
- 20 41. Hamilton-Wessler, M., Ader, M., Getty, L., Dea, M., Hamm-Alvarez, S., and Bergman, R.N. Microtubule-disrupting agent taxol reduces insulin-stimulated glucose uptake *in vivo*. *Diabetes*, 45 (Suppl. 2): Abstr. 615, 1996.
42. Brem, S. Angiogenesis antagonists: current clinical trials. *Angiogenesis*, 2: 9-20, 1998.
- 25 43. Harris, A.L. Are angiostatin and endostatin cures for cancer? *Lancet*, 351: 1598-1599, 1998.
44. Appel R and Berger G. Über das Hydrazidosulphamide (on hydrazidosulphamide) *Chemische Berichte* 1958; 91; 1339.
45. Perrin DD and Amarego WLF. Purification of laboratory chemicals. *Pergamon press; Oxford* 1988; 290.
- 30

CLAIMS

1. Use of a cyclic compound or a pharmaceutically active salt thereof in the manufacture of a medicament to prevent and/or inhibit and/or arrest cell cycling, wherein the cyclic compound comprises at least one ring,
wherein Group I and Group II, independently of each other, are attached to a ring of the cyclic compound;
wherein Group I is a hydrocarbyl or an oxyhydrocarbyl group; and
wherein Group II is a group of the formula



X is P or S;

when X is P, Y is =O or S, Z is -OH and R is hydrocarbyl or H;

when X is S, Y is =O, Z is =O, and R is hydrocarbyl or N(R₁)(R₂), wherein each of R₁ and R₂ is independently selected from H or a hydrocarbyl group.

2. Use of a compound according to claim 1 in the manufacture of a medicament to irreversibly prevent and/or inhibit and/or arrest cell cycling.
3. Use according to claim 1 or 2 wherein cell cycling is prevented and/or inhibited and/or arrested in the G₂/M phase.
4. Use according to any one of the preceding claims wherein the cyclic compound is a polycyclic compound.
5. Use according to claim 4 wherein Group I and Group II are each attached to the same ring of the polycyclic compound.
6. Use according to claim 4 or 5 the polycyclic compound has a steroidal structure.
7. Use according to any one of the preceding claims wherein Group I and Group II are

each attached to the same ring at positions *ortho* with respect to each other.

8. Use according to claim 6 or 7 wherein Group I is attached to the 2 position of the A ring of the steroidal nucleus.

5

9. Use according to claim 6, 7 or 8 wherein Group II is attached to the 3 position of the A ring of the steroidal nucleus.

10. Use according to any one of the preceding claims wherein Group I is an oxyhydrocarbyl group, preferably a group of the formula $C_{1-6}O$.

10

11. Use according to claim 10 wherein the group $C_{1-6}O$ is a methoxy group.

15

12. Use according to claim 11 wherein the compound is 2-methoxyoestrone-3-*O*-sulphamate.

13. Use according to any one of claims 1 to 9 wherein Group I is a hydrocarbyl group, preferably a group of the formula C_{1-6} .

20

14. Use according to claim 13 wherein the group C_{1-6} is an ethyl group

15. Use according to claim 11 or 14 wherein the methoxy group or ethyl group is attached to the 2 position of the A ring of a steroidal nucleus.

25

16. Use according to any one of the preceding claims wherein X is S and R is $N(R_1)(R_2)$.


17. Use according to claim 16 wherein the compound is 2-methoxyoestrone-3-*O*-sulphamate or 2-ethyloestrone-3-*O*-sulphamate

30

18. Use according to any one of claims 1 to 15 wherein X is P; and Y is =O.

19. Use according to any one of claims 1 to 15 wherein X is S; and R is hydrocarbyl.

20. A sulphonate or a phosphonate compound comprising a steroidal ring and a sulphonate or a phosphonate group of the formula:



wherein

5 X is P or S;

when X is P, Z is -OH, and R is hydrocarbyl or H;

when X is S, Z is =O; and R is hydrocarbyl; and

wherein the sulphonate or phosphonate group is attached to the 3 position of the A ring of the steroidal nucleus; and at least one hydrocarbon group, preferably C₁₋₆ alkyl, is attached to the

10 2 position of the A ring of a steroidal nucleus.

21. Use of a composition in the manufacture of a medicament to prevent and/or inhibit and/or arrest cell cycling, wherein the composition comprises

(i) a compound as defined in any one of claims 1 to 17; and

15 (ii) a pharmaceutically acceptable carrier, diluent, or excipient; and/or

(iii) a biological response modifier.


22. Use according to claim 21 wherein the biological response modifier is a cytokine.

20 23. Use according to claim 22 wherein the cytokine is tumour necrosis factor (TNF).

24. Use of a cyclic compound or a pharmaceutically active salt thereof in the manufacture of a medicament to prevent and/or inhibit and/or arrest cell cycling, wherein the cyclic compound comprises at least one ring, wherein Group I and Group II, independently of each other, are attached to a ring of the cyclic compound;

25 wherein Group I is a hydrocarbyl or an oxyhydrocarbyl group; and

wherein Group II is a group of the formula



wherein each of R₁ and R₂ is independently selected from H or a hydrocarbyl group.

25. A method of treatment comprising administering to a subject in need of treatment a
5 compound or a composition as defined in any one of the preceding claims in order to
prevent and/or inhibit and/or arrest cell cycling.

1 / 16





FIG.1

2 / 16



3 / 16




FIG. 3

SUBSTITUTE SHEET (RULE 26)




FIG. 4





FIG. 5

6 / 16



7 / 16




FIG.7

8 / 16






FIG.8

9 / 16



SUBSTITUTE SHEET (RULE 26)

10 / 16




FIG.10

11 / 16





FIG.11

12 / 16



Synthesis of 2-ethyloestrone 3-O-sulphamate (11); (i) AlCl₃/AcCl, CH₃NO₂, 5 h, (ii) HBr/AcOH, Δ 2h, (iii) H₂/Pd-C (10%), THF/EtOH, 50 psi, 24h, (iv) AlCl₃/NaI; DCM/CH₃CN; Δ 5h, (v) NaH (1.2 eq.)/ DMF, ClSO₂NH₂, N₂, 12 h and (vi) NaBH₄/THF/TPA, 16 h.

FIG. 12

13 / 16




FIG.13


48 hours

24 hours

control

14 / 16

FIG. 14



15 / 16





FIG. 15

SUBSTITUTE SHEET (RULE 26)

FIG.16



(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
9 November 2000 (09.11.2000)

PCT

(10) International Publication Number
WO 00/66095 A3

- (51) International Patent Classification⁷: A61K 31/565, A61P 35/00
- (74) Agents: ALCOCK, David et al.; D Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).
- (21) International Application Number: PCT/GB00/01661
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 28 April 2000 (28.04.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
9910166.9 30 April 1999 (30.04.1999) GB
60/139,520 16 June 1999 (16.06.1999) US
0002113.9 28 January 2000 (28.01.2000) GB
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): STERIX LIMITED [GB/GB]; Magdalen Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB).
- Published:
with international search report
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): REED, Michael, John [GB/GB]; 42 Wimborne Gardens, London W13 8BZ (GB). POTTER, Barry, Victor, Lloyd [GB/GB]; 95 Dovers Park, Bathford, Bath BA1 7UE (GB).
- (88) Date of publication of the international search report:
9 August 2001
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: USE OF ESTRONE DERIVATIVES AS ANTITUMOUR AGENTS



(57) Abstract: There is provided use of a cyclic compound or a pharmaceutically active salt thereof in the manufacture of a medicament to prevent and/or inhibit and/or arrest cell cycling, wherein the cyclic compound comprises at least one ring, wherein Group I and Group II, independently of each other, are attached to a ring of the cyclic compound; wherein Group I is a hydrocarbyl or an oxyhydrocarbyl group; and wherein Group II is a group of formula (I), X is P or S; when X is P, Y is =O or S, Z is -OH and R is hydrocarbyl or H; when X is S, Y is =O, Z is =O, and R is hydrocarbyl or N(R₁)(R₂), wherein each of R₁ and R₂ is independently selected from H or a hydrocarbyl group.

WO 00/66095 A3

INTERNATIONAL SEARCH REPORT

International Application No

PC1/GB 00/01661

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K31/565 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 00 76487 A (POTTER BARRY VICTOR LLOYD ; REED MICHAEL JOHN (GB); STERIX LTD (GB)) 21 December 2000 (2000-12-21) the whole document	1-25
X	KLAUBER NANCY ET AL: "Inhibition of angiogenesis and breast cancer in mice by the microtubule inhibitors 2-methoxyestradiol and taxol." CANCER RESEARCH, vol. 57, no. 1, 1997, pages 81-86, XP002162122 ISSN: 0008-5472 the whole document	1-25

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

5 March 2001

Date of mailing of the international search report

19/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Orviz Diaz, P

INTERNATIONAL SEARCH REPORT

International Application No

PC1/GB 00/01661

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 99 64013 A (POTTER BARRY VICTOR LLOYD ; REED MICHAEL JOHN (GB); STERIX LTD (GB)) 16 December 1999 (1999-12-16) the whole document ---	1-25
X	PUROHIT A ET AL: "THE DEVELOPMENT OF A-RING MODIFIED ANALOGUES OF OESTRONE-3-O- SULPHAMATE AS POTENT STEROID SULPHATASE INHIBITORS WITH REDUCED OESTROGENICITY" JOURNAL OF STEROID BIOCHEMISTRY AND MOLECULAR BIOLOGY, GB, ELSEVIER SCIENCE LTD., OXFORD, vol. 64, no. 5, 1998, pages 269-275, XP000852539 ISSN: 0960-0760 the whole document ---	1-25
P,X	PUROHIT A: "Inhibition of tumor necrosis factor α -stimulated aromatase activity by microtubule-stabilizing agents, paclitaxel and 2-methoxyestradiol" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 261, 1999, pages 214-217, XP002121930 ISSN: 0006-291X the whole document ---	1-25
X	WO 93 05063 A (IMPERIAL COLLEGE) 18 March 1993 (1993-03-18) the whole document ---	1-25
X	WO 93 05064 A (IMPERIAL COLLEGE) 18 March 1993 (1993-03-18) the whole document ---	1-25
P,X	WO 99 27935 A (POTTER BARRY VICTOR LLOYD ; REED MICHAEL JOHN (GB); UNIV BATH (GB);) 10 June 1999 (1999-06-10) the whole document ---	1-25
X	CUSHMAN M ET AL: "SYNTHESIS, ANTITUBULIN AND ANTIMITOTIC ACTIVITY, AND CYTOTOXICITY OF ANALOGS OF 2-METHOXYESTRADIOL, AN ENDOGENOUS MAMMALIAN METABOLITE OF ESTRADIOL THAT INHIBITS TUBULIN POLYMERIZATION BY BINDING TO THE COLCHICINE BINDING SITE" JOURNAL OF MEDICINAL CHEMISTRY, US, AMERICAN CHEMICAL SOCIETY, WASHINGTON, vol. 38, no. 12, 1995, pages 2041-2049, XP002055798 ISSN: 0022-2623 the whole document ---	1-25

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International Application No

PC1/GB 00/01661

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCHWARZ S ET AL: "Synthesis of estrogen sulfamates: Compounds with a novel endocrinological profile" STEROIDS: STRUCTURE, FUNCTION, AND REGULATION,US,ELSEVIER SCIENCE PUBLISHERS, NEW YORK, NY, vol. 61, no. 12, 1 December 1996 (1996-12-01), pages 710-717, XP004016631 ISSN: 0039-128X the whole document ---	1-25
X	WOO L W L ET AL: "ACTIVE SITE DIRECTED INHIBITION OF ESTRONE SULFATASE BY NONSTEROIDAL COUMARIN SULFAMATES" JOURNAL OF MEDICINAL CHEMISTRY,US,AMERICAN CHEMICAL SOCIETY, WASHINGTON, vol. 39, no. 7, 1996, pages 1349-1351, XP000652210 ISSN: 0022-2623 the whole document ---	1-25
X	WO 98 24802 A (POTTER BARRY VICTOR LLOYD ; REED MICHAEL JOHN (GB); IMPERIAL COLLEG) 11 June 1998 (1998-06-11) the whole document ---	1-25
X	WOO L W L ET AL: "Oestrone 3-O-(N-acetyl)sulphamate, a potential molecular probe of the active site of oestrone sulphatase" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS,GB,OXFORD, vol. 7, no. 24, 16 December 1997 (1997-12-16), pages 3075-3080, XP004136588 ISSN: 0960-894X the whole document ---	1-25
P,X	WO 99 27936 A (POTTER BARRY VICTOR LLOYD ; REED MICHAEL JOHN (GB); UNIV BATH (GB);) 10 June 1999 (1999-06-10) the whole document --- -/--	1-25

INTERNATIONAL SEARCH REPORT

International Application No

PC1/GB 00/01661

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	<p>PUROHIT A ET AL: "RECENT ADVANCES IN THE DEVELOPMENT OF STEROID SULPHATASE INHIBITORS"</p> <p>JOURNAL OF STEROID BIOCHEMISTRY AND MOLECULAR BIOLOGY, GB, ELSEVIER SCIENCE LTD., OXFORD,</p> <p>vol. 69, no. 1/06, 1999, pages 227-238, XP000852540</p> <p>ISSN: 0960-0760</p> <p>the whole document</p> <p>& PROCEEDINGS OF THE 10TH INTERNATIONAL CONGRESS ON HORMONAL STEROIDS, QUEBEC, 17-21 JUNE 1998,</p> <p>-----</p>	1-25

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

The independent claim 1 relates to an extremely large number of compounds. A complete search is not possible. The search had to be limited to those compounds supported by the description, i.e. those explicitly mentioned in claim 17. Furthermore, claim 1 refers to a mechanism of action, rather to a therapeutic indication. The search focused on the antitumour effect of the compounds.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCi/GB 00/01661

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0076487 A	21-12-2000	NONE	
WO 9964013 A	16-12-1999	AU 4280799 A	30-12-1999
WO 9305063 A	18-03-1993	AT 191221 T	15-04-2000
		AU 2494992 A	05-04-1993
		DE 69230857 D	04-05-2000
		DE 69230857 T	02-11-2000
		EP 0602123 A	22-06-1994
		JP 7501514 T	16-02-1995
		US 6017904 A	25-01-2000
		US 5604215 A	18-02-1997
		US 5861390 A	19-01-1999
WO 9305064 A	18-03-1993	AT 194843 T	15-08-2000
		AU 717116 B	16-03-2000
		AU 1007799 A	04-03-1999
		AU 668882 B	23-05-1996
		AU 2490592 A	05-04-1993
		AU 689043 B	19-03-1998
		AU 5837396 A	05-09-1996
		AU 7195298 A	13-08-1998
		BR 9206434 A	27-09-1994
		CA 2114630 A	18-03-1993
		CZ 9400440 A	18-01-1995
		DE 69231280 D	24-08-2000
		DE 69231280 T	30-11-2000
		EP 0641355 A	08-03-1995
		EP 0928609 A	14-07-1999
		EP 0921130 A	09-06-1999
		EP 0982032 A	01-03-2000
		ES 2148178 T	16-10-2000
		FI 940903 A	20-04-1994
		FI 20002259 A	13-10-2000
		FI 20002260 A	13-10-2000
		FI 20002261 A	13-10-2000
		HU 66097 A	28-09-1994
		JP 2000038341 A	08-02-2000
		JP 7501515 T	16-02-1995
		JP 2000355542 A	26-12-2000
		JP 2000355598 A	26-12-2000
		NO 940635 A	29-04-1994
		NO 995075 A	29-04-1994
		NO 995076 A	29-04-1994
		NO 2595 A	29-04-1994
		SK 24694 A	09-11-1994
		US 6011024 A	04-01-2000
		US 6159960 A	12-12-2000
		US 5616574 A	01-04-1997
		US 5830886 A	03-11-1998
WO 9927935 A	10-06-1999	GB 2331988 A	09-06-1999
		AU 1345699 A	16-06-1999
		EP 1051177 A	15-11-2000
WO 9824802 A	11-06-1998	AU 5402398 A	29-06-1998
		EP 0942919 A	22-09-1999

Form PCT/ISA/210 (patent family annex) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC1/GB 00/01661

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9927936 A	10-06-1999	GB 2331987 A	09-06-1999
		AU 1345899 A	16-06-1999
		EP 1051178 A	15-11-2000
